

Physician Survey to Reassess Effectiveness of Strattera Risk Minimisation Activities

First published: 23/08/2013

Last updated: 22/02/2024

Study

Finalised

Administrative details

EU PAS number

EUPAS4577

Study ID

8929

DARWIN EU® study

No

Study countries

 Denmark

 Netherlands

 Sweden

 United Kingdom

Study description

This is a survey to re-assess the effectiveness of the atomoxetine risk minimisation activities among physicians who prescribe Strattera and/or monitor patients being treated with Strattera.

Study status

Finalised

Research institutions and networks

Institutions

GfK Health

First published: 01/02/2024

Last updated: 01/02/2024

Institution

Contact details

Study institution contact

Nicole Kellier nkellier@lilly.com

Study contact

nkellier@lilly.com

Primary lead investigator

Nicole Kellier

Primary lead investigator

Study timelines

Date when funding contract was signed

Actual: 26/04/2013

Study start date

Planned: 15/09/2013

Actual: 13/09/2013

Data analysis start date

Planned: 01/12/2013

Actual: 01/12/2013

Date of final study report

Planned: 28/03/2014

Actual: 27/03/2014

Sources of funding

- Pharmaceutical company and other private sector

More details on funding

Eli Lilly and Company

Study protocol

[RMiP PASS protocol_p1.pdf](#) (124.46 KB)

[RMiP PASS protocol_p1_mod.pdf](#) (180.31 KB)

Regulatory

Was the study required by a regulatory body?

No

Methodological aspects

Study type

Study type list

Study topic:

Human medicinal product

Study type:

Not applicable

Scope of the study:

Assessment of risk minimisation measure implementation or effectiveness

Data collection methods:

Primary data collection

Main study objective:

Re-assessment of the effectiveness of the atomoxetine risk minimisation activities among physicians who prescribe Strattera and/or monitor patients being treated with Strattera. The goal of these activities is to inform HCPs of the cardiovascular or cerebrovascular disorder contraindication and SPC recommendation to monitor BP and HR in all patients at baseline and during treatment with Strattera

Study drug and medical condition

Anatomical Therapeutic Chemical (ATC) code

(N06BA09) atomoxetine

atomoxetine

Population studied

Short description of the study population

Physicians (Paediatricians, Child/Adolescent Psychiatrists or other NonPaediatricians and GPs) who prescribe Strattera and/or monitor patients treated with Strattera

Age groups

- Adults (18 to < 46 years)
 - Adults (46 to < 65 years)
 - Adults (65 to < 75 years)
 - Adults (75 to < 85 years)
 - Adults (85 years and over)
-

Estimated number of subjects

550

Study design details

Outcomes

The risk minimisation activities assessed by this survey include sustained knowledge and awareness of the risk messages, The survey will also include an assessment of awareness and adherence to the changes in the SmPC requirements specific to cardiovascular risks and monitoring

Data analysis plan

Data analyses will be descriptive and will entail tabular displays of mean values and the frequency distribution of item responses.

Documents

Study results

[B024 PASS.pdf](#) (1.87 MB)

Study, other information

[RMiP PASS protocol_p2.pdf](#) (276.74 KB)

Data management

ENCePP Seal

The use of the ENCePP Seal has been discontinued since February 2025. The ENCePP Seal fields are retained in the display mode for transparency but are no longer maintained.

Data sources

Data sources (types)

[Other](#)

Data sources (types), other

Survey

Use of a Common Data Model (CDM)

CDM mapping

No

Data quality specifications

Check conformance

Unknown

Check completeness

Unknown

Check stability

Unknown

Check logical consistency

Unknown

Data characterisation

Data characterisation conducted

No