Longitudinal Analyses of Body Mass Index and Risk of Parkinson's Disease in 2 million people over 2 decades (BMI and Parkinson's disease)

First published: 04/04/2018 Last updated: 02/07/2024





Administrative details

EU PAS number
EUPAS23454
Study ID
23455
DARWIN EU® study
No
Study countries
Spain
United Kingdom

Study description

Previous research suggests that people who are overweight or obese may have a higher risk of developing Parkinson's disease than people with normal weight, while other studies show no such association. The objective of this study is therefore to investigate the association between BMI and risk of Parkinson's disease. A cohort will be derived from CPRD of people aged 40 years or older with a first BMI recording between 1992 and 2007. People with a prior record of Parkinson's disease or dementia will be excluded. Incidence rates of Parkinson's disease will be calculated for each BMI category using Poisson regression, adjusting for differences in patient characteristics and allowing for competing risks with a novel approach. This study will provide information from a very large number of people (2 million) with a sizeable amount of follow-up data in a representative sample. The findings will therefore provide important information to help clarify the relationship between BMI and Parkinson's disease to inform preventative and therapeutic strategies for Parkinson's disease.


Study status

Ongoing

Research institutions and networks

Institutions

OXON Epidemiology
Spain
United Kingdom
First published: 06/12/2010
Last updated: 15/03/2024



Contact details

Study institution contact

Stuart Pocock Stuart.Pocock@lshtm.ac.uk

Study contact

Stuart.Pocock@lshtm.ac.uk

Primary lead investigator

Nawab Qizilbash MBChB MRCP(UK) BSc MSc DPhil(Oxon.)

Primary lead investigator

Study timelines

Date when funding contract was signed

Planned: 20/07/2016

Actual: 20/07/2016

Study start date

Planned: 20/07/2016

Actual: 20/07/2016

Data analysis start date

Planned: 15/09/2016

Actual: 15/09/2016

Date of final study report

Planned: 17/09/2018

Sources of funding

Other

More details on funding

No funding

Regulatory

Was the study required by a regulatory body?

No

Is the study required by a Risk Management Plan (RMP)?

Not applicable

Methodological aspects

Study type

Study type list

Study type:

Non-interventional study

Scope of the study:

Assessment of risk minimisation measure implementation or effectiveness Disease epidemiology

Main study objective:

The primary objective is to estimate the age-specific associations between BMI and Parkinson's disease. Secondary objectives are to assess the shape of the association and Identify potential modifiers and confounders of the risk. The results will be usable for disease prediction models, patient management, the planning of trials, and help in hypotheses of pathophysiological mechanisms of PD

Study Design

Non-interventional study design

Cohort

Study drug and medical condition

Medical condition to be studied

Obesity

Parkinson's disease

Population studied

Short description of the study population

2 million patients followed up over 2 decades. Underweight (< 20 kg/m2): 106,716 people Healthy weight (20-24.9 kg/m2):629,126 people Overweight (25-29.9 kg/m2): 727,339 people Obese (\geq 30 kg/m2): 489,406 people With a median follow-up of 9·3 years, PD occurred in 11,616 people.

Age groups

Adults (46 to < 65 years)

Adults (65 to < 75 years)

Adults (75 to < 85 years)

Adults (85 years and over)

Estimated number of subjects

2000000

Study design details

Outcomes

Parkinson's disease

Data analysis plan

We compared rates of PD across categories of BMI using Poisson regression models. Standardised incidence rates and rate ratios were either adjusted for age (in five-year bands) and sex, or further adjusted for the following covariates, measured at the time of index BMI. We used age at diagnosis, updating the data as people moved through the age categories during follow-up. Incidence rates were standardised to the age and sex distribution of the overall study population. We performed analyses to investigate whether the association between BMI and PD varied depending on the time from the index BMI measurement. We calculated the cumulative incidence of PD using the Kaplan-Meier estimator, with age as the underlying time scale. We performed a sensitivity analysis to investigate whether our findings could be reasonably explained by competing risks using a novel approach under a hypothetical scenario and using matching

Documents

Study, other information

BMI and PD poster ICPE 2016.pdf(465.51 KB)

Study publications

Iwagami, M., Qizilbash, N., Gregson, J., Douglas, I., Johnson, M., Pearce, N., ...

Data management

ENCePP Seal

The use of the ENCePP Seal has been discontinued since February 2025.

The ENCePP Seal fields are retained in the display mode for transparency but are no longer maintained.

Data sources

Data source(s)

Clinical Practice Research Datalink

Data sources (types)

Electronic healthcare records (EHR)

Use of a Common Data Model (CDM)

CDM mapping

No

Data quality specifications

Unknown Check completeness Unknown

Check stability

Check conformance

Unknown

Check logical consistency

Unknown

Data characterisation

Data characterisation conducted

No