

# Drug Utilization Study for Pirinase Hayfever Relief for Adults 0.05% Nasal Spray (205708)

**First published:** 12/02/2016

**Last updated:** 01/04/2024

Study

Finalised

## Administrative details

### EU PAS number

EUPAS12436

---

### Study ID

25056

---

### DARWIN EU® study

No

---

### Study countries

 United Kingdom

---

### Study description

The Medicines and Healthcare products Regulatory Agency (MHRA) has required this study to evaluate consumer compliance with the product labelling of Pirinase Hayfever Relief for Adults 0.05% Nasal Spray over at least 2 allergy seasons. The purpose is to obtain real-world information on how consumers are complying with the product labelling This study will coincide with the launch of Pirinase Hayfever Relief for Adults 0.05% Nasal Spray in the United Kingdom (UK).The 5 primary objectives are:1. To assess if consumers of the correct age use the product: Ages 18 and older2. To assess the correct frequency of use: No more than 2 sprays in each nostril per day3. To assess reduction of dose: If symptoms improve, 1 spray in each nostril per day4. To assess if a physician is consulted before use: if a woman is pregnant or breastfeeding5. To assess if a physician is consulted: If symptoms have not improved after using for 7 days, a doctor is consultedThe 2 secondary objectives are:1. To assess if consumers do not use if they are taking medications for Human Immunodeficiency Virus (HIV)2. To assess if a physician is consulted: Not used more than 1 month continuously without consulting a doctor

---

### **Study status**

Finalised

## Research institutions and networks

### Institutions

[Concentrics Research](#)

## Contact details

**Study institution contact**

GSK Clinical Disclosure Advisor GSK Clinical Disclosure  
Advisor Cdr\_mailbox@gsk.com

Study contact

[Cdr\\_mailbox@gsk.com](mailto:Cdr_mailbox@gsk.com)

**Primary lead investigator**

GSK Clinical Disclosure Advisor GSK Clinical Disclosure  
Advisor

Primary lead investigator

## Study timelines

**Date when funding contract was signed**

Planned: 15/02/2016

Actual: 29/01/2016

---

**Study start date**

Planned: 15/02/2016

Actual: 04/04/2016

---

**Date of final study report**

Planned: 30/09/2018

Actual: 19/07/2018

## Sources of funding

- Pharmaceutical company and other private sector

## More details on funding

GlaxoSmithKline

## Study protocol

[gsk-205708-protocol-redact.pdf](#) (1.37 MB)

## Regulatory

### **Was the study required by a regulatory body?**

Yes

---

### **Is the study required by a Risk Management Plan (RMP)?**

EU RMP category 3 (required)

## Methodological aspects

### Study type

### Study type list

#### **Study topic:**

Disease /health condition

Human medicinal product

---

**Study type:**

Not applicable

---

**Main study objective:**

The primary objective of this study is to evaluate if consumers comply with key warnings and directions on the outer label for selection and use of the drug.

## Study drug and medical condition

**Medicinal product name, other**

Fluticasone propionate

---

**Medical condition to be studied**

Rhinitis allergic

## Population studied

**Short description of the study population**

Approximately 1537 consumers in the UK who have purchased and used Pirinase Hayfever Relief for Adults 0.05% Nasal Spray for at least 7 days.

---

**Age groups**

- Children (2 to < 12 years)
- Adolescents (12 to < 18 years)
- Adults (18 to < 46 years)
- Adults (46 to < 65 years)
- Adults (65 to < 75 years)

- Adults (75 to < 85 years)
  - Adults (85 years and over)
- 

### **Special population of interest**

Hepatic impaired

Immunocompromised

Pregnant women

Renal impaired

---

### **Estimated number of subjects**

1537

## Study design details

### **Outcomes**

To assess:- if consumers of the correct age use the product: (18 and older)- the correct frequency of use: No more than 2 sprays in each nostril/day- reduction of dose: If symptoms improve, 1 spray in each nostril/day- if a physician is consulted before use: if a woman is pregnant or breastfeeding- if a physician is consulted: If symptoms have not improved after using for 7 days, The two secondary objectives are the following:1. To assess if consumers do not use if they are taking medications for Human Immunodeficiency Virus (HIV)2. To assess if a physician is consulted: Not used more than 1 month continuously without consulting a doctor

---

### **Data analysis plan**

Primary endpoints will be analyzed individually and no adjustment for multiple comparisons will be performed. The compliance rate for each of the five primary endpoints will be reported and its two-sided 95% confidence limit will be computed using the exact method. The endpoint will have met the threshold for

success if the lower bound of the 2-sided 95% exact confidence interval (CI) is at least 80%. Secondary endpoints will be analyzed individually and no adjustment for multiple comparisons will be performed. The compliance rate for each of the secondary endpoints will be reported and its two-sided 95% confidence limit will be computed using the exact method. There are no success thresholds set for secondary endpoints.

## Data management

### ENCePP Seal

The use of the ENCePP Seal has been discontinued since February 2025. The ENCePP Seal fields are retained in the display mode for transparency but are no longer maintained.

## Data sources

### Data sources (types)

[Other](#)

---

### Data sources (types), other

Information direct from consumers in the UK who have purchased and used the product

## Use of a Common Data Model (CDM)

### CDM mapping

No

## Data quality specifications

### **Check conformance**

Unknown

---

### **Check completeness**

Unknown

---

### **Check stability**

Unknown

---

### **Check logical consistency**

Unknown

## Data characterisation

### **Data characterisation conducted**

Unknown