

# An Observational Post-Authorization Safety Specialist Cohort Event Monitoring Study (SCEM) to Monitor the Safety and Utilization of Rivaroxaban (Xarelto®) for the Prevention of Stroke in Patients with AF, Treatment of DVT and PE, and the Prevention of Recurrent DVT and PE in the Secondary Care Hospital Setting in England and Wales (ROSE)

**First published:** 22/05/2013

**Last updated:** 01/04/2024

Study

Finalised

## Administrative details

### EU PAS number

EUPAS3979

---

### Study ID

28398

---

## **DARWIN EU® study**

No

---

### **Study countries**

☐ United Kingdom

---

### **Study description**

A study to evaluate the use and short term safety of Rivaroxaban (Xarelto®) during real-life use in Secondary Care (hospitals) in England and Wales.

Rivaroxaban is an anti-coagulant medication which has previously been used only in patients having hip or knee replacement surgery but will now also be used to prevent stroke and systemic embolism in patients with non-valvular atrial fibrillation. Rivaroxaban will also be used to treat deep vein thrombosis (DVT) and pulmonary embolism (PE) and to prevent recurrent DVT and PE. This study aims to evaluate the use of rivaroxaban and its short term safety when used by patients for these new indications. This study was requested by the Committee for Medicinal products for Human use (CHMP). The study will recruit patients starting rivaroxaban treatment and asking their care team to answer some simple questions about them at the time they start and again in 12 weeks time. The study will also recruit patients starting alternative anticoagulant therapies and their care team will be asked the same questions. These patients will be used to compare the differences between users of rivaroxaban and users of alternative anticoagulant therapies. If a participant has an adverse event during the 12 week period, we may ask the patient's care team to fill out a further follow up questionnaire. No other examinations or tests will be performed. The participant's consent will be obtained to access their medical charts. Any adult patient started by their care team on rivaroxaban or an alternative anticoagulant for the specified indications during the study period will be eligible to take part. It is a national study covering the whole of England and Wales. The study will last for approximately 3 years of data collection (in

order to reach a cohort of 3400 patients - 1700 rivaroxaban users and 1700 alternative anticoagulant therapy users), although each individual patient will only be involved for a 12 week period of observation.

---

## Study status

Finalised

## Research institutions and networks

### Institutions

#### Drug Safety Research Unit (DSRU)

☐ United Kingdom

**First published:** 10/11/2021

**Last updated:** 16/02/2024

Institution

Not-for-profit

ENCePP partner

### Networks

Stroke Research Network, Non-malignant  
Haematology Group, Cardiovascular Group

### Contact details

**Study institution contact**

Alison Evans [alison.evans@dsru.org](mailto:alison.evans@dsru.org)

Study contact

[alison.evans@dsru.org](mailto:alison.evans@dsru.org)

**Primary lead investigator**

Saad Shakir

Primary lead investigator

## Study timelines

**Date when funding contract was signed**

Planned: 19/12/2012

Actual: 19/12/2012

---

**Study start date**

Planned: 21/05/2013

Actual: 05/09/2013

---

**Data analysis start date**

Planned: 01/11/2016

Actual: 31/03/2017

---

**Date of interim report, if expected**

Planned: 13/04/2015

Actual: 13/08/2015

---

**Date of final study report**

Planned: 28/02/2017

Actual: 26/06/2017

## Sources of funding

- Pharmaceutical company and other private sector

## More details on funding

Bayer plc

## Study protocol

[ROSE protocol\\_v40\\_20\\_11\\_2014.pdf](#)(1.25 MB)

## Regulatory

**Was the study required by a regulatory body?**

Yes

---

**Is the study required by a Risk Management Plan (RMP)?**

EU RMP category 3 (required)

## Methodological aspects

### Study type

### Study type list

**Study topic:**

Disease /health condition  
Human medicinal product

---

**Study type:**

Non-interventional study

---

**Scope of the study:**

Assessment of risk minimisation measure implementation or effectiveness  
Drug utilisation

**Data collection methods:**

Primary data collection

---

**Main study objective:**

To monitor the short-term (up to 12 weeks) safety profile and drug utilisation of rivaroxaban as prescribed to patients for medical conditions requiring anticoagulation by specialist Healthcare Professionals in the secondary care hospital setting in England and Wales.

## Study Design

**Non-interventional study design**

Cohort

Other

---

**Non-interventional study design, other**

Intensive monitoring schemes

## Study drug and medical condition

**Name of medicine**

XARELTO

---

**Medical condition to be studied**

Atrial fibrillation

Deep vein thrombosis

Pulmonary embolism

## Population studied

**Short description of the study population**

Rivaroxaban prescribed newuser adult patients (i.e. rivaroxaban naïve who may or may not be antithrombotic therapy naïve) for the prevention of stroke and systemic embolism in adult patients with non-valvular atrial fibrillation, the treatment of deep vein thrombosis (DVT), pulmonary embolism (PE), and the prevention of recurrent DVT and PE in adult patients, requiring anticoagulation under normal conditions of use in the secondary care hospital setting.

---

**Age groups**

Adults (18 to < 46 years)

Adults (46 to < 65 years)

Adults (65 to < 75 years)

Adults (75 to < 85 years)

Adults (85 years and over)

---

**Estimated number of subjects**

3400

## Study design details

## Outcomes

To provide timely information for rivaroxaban users on the estimation of the cumulative incidence of bleeding events

1.Prescriber/cohort accrual, type of prescriber, setting 2.Non-clinical reasons for prescribing, prognostic health factors, clinical risk factors for haemorrhage 3.Changes of health profile/adherence/treatment 4.Quantify risk of haemorrhage within GI/urogenital/intracranial sites for contextual cohort & within critical organ sites for both cohorts, major & non-major bleeds, other events.

---

## Data analysis plan

Data analysis will include • Response rates to describe recruitment • Hazard rates to explore the incidence of selected events • Descriptive analyses of baseline health profile of patients • Analysis of risk and incidence densities to describe the risk profile of events reported in patient subgroups of special interest

# Documents

## Study results

[EU-PAS\\_Abstract.pdf](#)(458.46 KB)

---

## Study report

[CSR 16171 R-11983\\_EUPAS.pdf](#)(6.13 MB)

## Data management

## ENCePP Seal



The use of the ENCePP Seal has been discontinued since February 2025. The ENCePP Seal fields are retained in the display mode for transparency but are no longer maintained.

## Data sources

### Data sources (types)

Other

---

### Data sources (types), other

Prospective patient-based data collection

## Use of a Common Data Model (CDM)

### CDM mapping

No

## Data quality specifications

### Check conformance

Unknown

---

### Check completeness

Unknown

---

### Check stability

Unknown

---

**Check logical consistency**

Unknown

Data characterisation

**Data characterisation conducted**

Unknown