

Incidence of tolerance, dependence, drug abuse and misuse of Targin in patients with Restless Legs Syndrome

First published: 17/05/2021

Last updated: 26/03/2024

Study

Finalised

Administrative details

PURI

<https://redirect.ema.europa.eu/resource/41661>

EU PAS number

EUPAS41097

Study ID

41661

DARWIN EU® study

No

Study countries

Germany

Study status

Finalised

Research institutions and networks

Institutions

Institute for Applied Health Research Berlin (InGef)

Germany

First published: 01/02/2024

Last updated: 01/02/2024

Institution

Educational Institution

Contact details

Study institution contact

Lee Seuunghee

Study contact

SeungHee.Lee@mundipharma.co.kr

Primary lead investigator

Lee Seuunghee

Primary lead investigator

Study timelines

Date when funding contract was signed

Planned: 28/02/2018

Actual: 28/02/2018

Study start date

Planned: 01/10/2020

Actual: 01/10/2020

Date of final study report

Planned: 29/03/2021

Actual: 29/03/2021

Sources of funding

- Pharmaceutical company and other private sector

More details on funding

Mundipharma

Regulatory

Was the study required by a regulatory body?

Unknown

Is the study required by a Risk Management Plan (RMP)?

Not applicable

Methodological aspects

Study type

Study type list

Study topic:

Disease /health condition
Human medicinal product

Study type:

Non-interventional study

Scope of the study:

Assessment of risk minimisation measure implementation or effectiveness

Data collection methods:

Secondary use of data

Main study objective:

To estimate the incidence of problematic Targin prescription use and abuse in patients treated with Targin for RLS.

Study Design

Non-interventional study design

Cohort
Other

Non-interventional study design, other

Prescription event monitoring

Study drug and medical condition

Name of medicine, other

Targin

Medical condition to be studied

Restless legs syndrome

Population studied

Short description of the study population

Patients with Restless Legs Syndrome.

Age groups

Adults (18 to < 46 years)

Adults (46 to < 65 years)

Adults (65 to < 75 years)

Adults (75 to < 85 years)

Adults (85 years and over)

Estimated number of subjects

2757

Study design details

Outcomes

confirmed problematic Targin use had a low incidence rate and high doses and long prescribing were also reasonably uncommon while there was a moderate proportion of patients using other opioids during Targin exposure. While confirmed opioid abuse/dependency may be underreported in a routine

healthcare database such as InGef, not all of the identified occurrences may be attributed to Targin. To characterise Targin for RLS treatment patterns in patients To estimate the incidence of “suspected doctor shopping” in patients treated with Targin for RLS To estimate the amount of off-label use of Targin in RLS patients To determine concomittant treatment in patients treated with Targin for RLS. To estimate incidences of pre-specified adverse events during treatment

Data analysis plan

Statistical analysis were carried out using the data environment used by InGef.

Data management

Data sources

Data sources (types)

[Drug registry](#)

[Other](#)

Data sources (types), other

Prescription event monitoring

Use of a Common Data Model (CDM)

CDM mapping

No

Data quality specifications

Check conformance

Unknown

Check completeness

Unknown

Check stability

Unknown

Check logical consistency

Unknown

Data characterisation

Data characterisation conducted

No