# US-based cross-sectional survey in patients taking lasmiditan: Functional REstoration with rEyvow (FREE)

First published: 17/11/2020

Last updated: 13/12/2024





# Administrative details

PURI
https://redirect.ema.europa.eu/resource/41778
EU PAS number
EUPAS37245
Study ID
41778
DARWIN EU® study
No
Study countries
United States

#### **Study description**

The objective of the study is to assess patients' ability to return to their usual daily activities after treating a migraine attack with lasmiditan. Study data will be collected by conducting a 15-minute web-based cross-sectional survey in adult patients in US who have redeemed a REYVOW savings card and have taken lasmiditan at least once in the past month for treatment of a migraine attack.

#### **Study status**

Finalised

## Research institutions and networks

#### **Institutions**



## Contact details

**Study institution contact**Louise Lombard

Study contact

#### lombard louise@lilly.com

#### **Primary lead investigator**

#### **Amit Qamra**

**Primary lead investigator** 

# Study timelines

#### Date when funding contract was signed

Actual: 08/07/2020

#### Study start date

Actual: 16/11/2020

#### Data analysis start date

Planned: 01/12/2020

Actual: 01/12/2020

#### **Date of final study report**

Planned: 01/06/2021

Actual: 15/06/2021

# Sources of funding

• Pharmaceutical company and other private sector

## More details on funding

Eli Lilly and Company

# Study protocol

# Regulatory

Was the study required by a regulatory body?

No

Is the study required by a Risk Management Plan (RMP)?

Not applicable

# Methodological aspects

# Study type

# Study type list

#### **Study topic:**

Disease /health condition

Human medicinal product

#### Study type:

Non-interventional study

#### Scope of the study:

Effectiveness study (incl. comparative)

#### **Data collection methods:**

#### Main study objective:

The objective of the study is to assess patients' ability to return to their usual daily activities after treating a migraine attack with lasmiditan.

## Study Design

#### Non-interventional study design

Cross-sectional

# Study drug and medical condition

**Study drug International non-proprietary name (INN) or common name**LASMIDITAN

#### Medical condition to be studied

Migraine

# Population studied

#### Short description of the study population

Adult patients in US who have redeemed a REYVOW savings card and have taken lasmiditan at least once in the past month for treatment of a migraine attack prior to completing the survey.

Inclusion Criteria

The study will include all participants who meet the following criteria:

- 1. Age ≥18 at time of study
- 2. Enrolled in US REYVOW patient support program
- 3. Redeemed a REYVOW Savings Card
- 4. Provide informed consent to participate in the study
- 5. Have taken lasmiditan in the past month prior to completing the survey Exclusion Criteria

Participants will be excluded from the study:

- 1. Cannot read or understand English
- 2. Cannot use a computer or smart device with internet access

#### Age groups

Adults (18 to < 46 years)

Adults (46 to < 65 years)

Adults (65 to < 75 years)

Adults (75 to < 85 years)

Adults (85 years and over)

#### **Special population of interest**

Other

#### Special population of interest, other

Migrane patients

#### **Estimated number of subjects**

250

# Study design details

#### **Outcomes**

Ability to perform and the time taken to return to usual daily activities engaged in when the migraine attack started, and Ability to return to planned activities for that day and level of impairment of those activities, Activity that was most impacted by their migraine attack prior to taking lasmiditan, the activity that the patient was most concerned about prior to taking lasmiditan, and the level of impairment of usual activities during an untreated or unsuccessfully treated attack prior to when lasmiditan was initiated

#### Data analysis plan

All study measures will be reported descriptively using means, standard deviations, medians, and 95% confidence intervals for continuous outcome variables and frequencies and percentages for categorical outcome variables. Differences between subgroups will be evaluated: analyses will be conducted to examine whether the select study outcome measures differ by each of the subgroups, using chi-square tests for categorical variables and one-way analysis-of-variance tests (ANOVAs) for continuous variables. P-values will be provided for the omnibus test (one-way ANOVA) and pair-wise testing between the groups. It should be noted that all comparisons are conditional on whether each subgroup results in adequate sample size.

### **Documents**

#### **Study results**

REYVOW FREE Report\_ENCePP version. 14 JUN 2021\_APPROVED.pdf(5.17 MB)

#### Study publications

Schwedt TJ, Lombard LA, Doty E, Vincent M, Mills KM, Ayer DW, et al. Functional...

## Data management

#### Data sources

Data sources (types) Other
Data sources (types), other Electronic one-time survey
Use of a Common Data Model (CDM)
CDM mapping No
Data quality specifications
Check conformance
Unknown
Check completeness
Unknown
Check stability
Unknown
Check logical consistency
Unknown
Data characterisation
Data characterisation conducted

No