

# A Prospective cohort study on the use of Sofosbuvir-based regimens in HCV-infected patients in clinical practice in France (HELIOS)

**First published:** 24/09/2015

**Last updated:** 22/02/2024

Study

Finalised

## Administrative details

### EU PAS number

EUPAS11074

### Study ID

50593

### DARWIN EU® study

No

### Study countries

☐ France

### Study description

GS-FR-334-1530: This was a multi-centre, prospective, non interventional cohort study. This study collected and evaluated information on safety and efficacy of sofosbuvir-based regimens in routine clinical practice in France.

---

## Study status

Finalised

## Research institutions and networks

### Institutions

#### Gilead Sciences

**First published:** 12/02/2024

**Last updated:** 12/02/2024

Institution

Pharmaceutical company

Multiple centres: 45 centres are involved in the study

## Contact details

### Study institution contact

Gilead Study Director [ClinicalTrialDisclosure@gilead.com](mailto:ClinicalTrialDisclosure@gilead.com)

Study contact

**Primary lead investigator**  
Gilead Study Director

Primary lead investigator

## Study timelines

### **Date when funding contract was signed**

Planned: 30/09/2014

Actual: 30/09/2014

---

### **Study start date**

Planned: 05/10/2015

Actual: 13/10/2015

---

### **Data analysis start date**

Planned: 30/11/2018

Actual: 23/04/2021

---

### **Date of final study report**

Planned: 31/12/2021

Actual: 26/10/2022

## Sources of funding

- Pharmaceutical company and other private sector

## More details on funding

## Study protocol

[GS-FR-334-1530 Protocol-5Dec2014 - Final.pdf](#) (3.01 MB)

[amd-2-prot-GS-FR-334-1530-09aug2017.pdf](#) (2.27 MB)

## Regulatory

**Was the study required by a regulatory body?**

No

---

**Is the study required by a Risk Management Plan (RMP)?**

Not applicable

## Methodological aspects

### Study type

### Study type list

**Study topic:**

Disease /health condition

Human medicinal product

---

**Study type:**

Non-interventional study

---

**Scope of the study:**

Effectiveness study (incl. comparative)

Safety study (incl. comparative)

**Data collection methods:**

Primary data collection

---

**Main study objective:**

The primary objective of this study was to assess the efficacy of sofosbuvir-based regimens in adult patients with chronic hepatitis C virus infection treated in routine clinical practice.

## Study Design

**Non-interventional study design**

Cohort

Other

---

**Non-interventional study design, other**

Multi-centre, prospective, study

## Study drug and medical condition

**Medical condition to be studied**

Chronic hepatitis C

## Population studied

## **Short description of the study population**

Adults patients aged 18 years or older with hepatitis C virus (HCV) infection received treatment with sofosbuvir, ledipasvir/sofosbuvir or sofosbuvir/velpatasvir in clinical practice in France.

Inclusion Criteria:

- ☐ Age  $\geq$  18 years
- ☐ HCV-infected patients (HCV RNA positive)
- ☐ Patient initiating treatment with Sofosbuvir\*, Ledipasvir/Sofosbuvir\* or Sofosbuvir/Velpatasvir\*
- ☐ Signed Patient Information Form

\* prescribed according to the respective SmPC

Exclusion Criteria:

- ☐ Concurrent participation in a HCV clinical trial (except trials not testing investigational medicinal product)
- ☐ Patients presenting a risk of not being able to be followed during 2 years (patients planning to move home, or leave the country in a foreseeable future)

---

## **Age groups**

Adults (18 to < 46 years)

Adults (46 to < 65 years)

Adults (65 to < 75 years)

Adults (75 to < 85 years)

Adults (85 years and over)

---

## **Special population of interest**

Other

---

## **Special population of interest, other**

## **Estimated number of subjects**

1031

## Study design details

### **Outcomes**

The proportion of subjects with sustained virologic response 12 weeks after discontinuation of therapy (SVR12). Rates of Adverse Drug Reaction, proportion of subjects with virologic response at 24 weeks post treatment, quality of life, patient adherence to HCV treatment, work productivity and activity Impairment, rate of consultations/hospitalizations due to HCV and/or liver disease and/or HCV treatment, treatment occurrence for anemia, thrombopenia and/or neutropenia

---

### **Data analysis plan**

Descriptive analysis Binary, categorical and ordinal variables were described by counts and frequencies of each modality (over the total number of responses), and comparisons between groups were tested using chi square tests and/or Fisher's exact tests. Continuous variables were described by means, standard errors, 95% confidence intervals, medians, minima, and maxima. Differences in means and medians between groups were tested using t-tests and non-parametric tests, respectively. When deemed necessary, sub-group comparisons and/or between time point comparisons may have been implemented. Multivariable analysis As part of the secondary objectives of this study, rates of events per person-time of exposure were computed using Poisson regression, after adjusting for potential confounding factors. These events included medication switching, medication discontinuation, co-medication use, ADRs and SADR and fatal AEs.

## Documents

## Study results

[GS-FR-334-1530-CSR synopsis\\_f-redact.pdf](#) (174.97 KB)

---

## Data management

### ENCePP Seal

The use of the ENCePP Seal has been discontinued since February 2025. The ENCePP Seal fields are retained in the display mode for transparency but are no longer maintained.

## Data sources

### Data sources (types)

[Other](#)

---

### Data sources (types), other

Prospective patient-based data collection

## Use of a Common Data Model (CDM)

### CDM mapping

No

## Data quality specifications



**Check conformance**

Unknown

---

**Check completeness**

Unknown

---

**Check stability**

Unknown

---

**Check logical consistency**

Unknown

## Data characterisation

**Data characterisation conducted**

No