

A Retrospective Chart Review Study of Patients With Chronic Lung Allograft Dysfunction-Bronchiolitis Obliterans Syndrome (CLAD-BOS) Post Lung Transplantation – "BOSTON EVOLUTION"

First published: 16/06/2026

Last updated: 16/06/2026

Study

Finalised

Administrative details

EU PAS number

EUPAS1000001016

Study ID

1000001016

DARWIN EU® study

No

Study countries

 Belgium

 Spain

Study description

A retrospective, multinational, multicenter chart review study of adult patients with clinically diagnosed Chronic Lung Allograft Dysfunction-Bronchiolitis Obliterans Syndrome (CLAD-BOS) after lung transplantation. All data retrospectively collected from patient records starting from lung transplantation date through last date of available data.

Study status


Finalised

Research institutions and networks

Institutions

Fortrea - Real world Intelligence & Late Phase Solution

 Germany

 United Kingdom (Northern Ireland)

First published: 15/12/2015

Last updated: 31/10/2023

Institution

Non-Pharmaceutical company

ENCePP partner

Zambon

First published: 01/02/2024

Last updated: 01/02/2024

Institution

Contact details

Study institution contact

Paola Castellani clinicaltrials@zambongroup.com

Study contact

clinicaltrials@zambongroup.com

Primary lead investigator

Paola Castellani

Primary lead investigator

Study timelines

Date when funding contract was signed

Planned: 30/11/2023

Actual: 30/11/2023

Study start date

Planned: 18/05/2024

Actual: 27/06/2024

Data analysis start date

Planned: 14/01/2025

Actual: 12/05/2025

Date of final study report

Planned: 15/04/2025

Actual: 16/04/2026

Sources of funding

- Pharmaceutical company and other private sector

More details on funding

Zambon SpA, Via Lillo del Duca, 10 Bresso (MI) 20091 Italy

Regulatory

Was the study required by a regulatory body?

No

Is the study required by a Risk Management Plan (RMP)?

Not applicable

Other study registration identification numbers and links

Protocol version identifier: Z8000N01

Methodological aspects

Study type

Study type list

Study topic:

Disease /health condition

Study type:

Non-interventional study

Scope of the study:

Disease epidemiology

Other

If 'other', further details on the scope of the study

Study to describe natural disease evolution and clinical outcomes. Data collection methods: The study data sources consisted of medical records (including electronic and/or paper) and patient notes for the study patients.

Data collection methods:

Secondary use of data

Study design:

This was a retrospective, multinational, multicenter chart review study of adult patients with clinically diagnosed CLAD BOS after lung transplantation. All data were retrospectively collected from patient records, starting from the lung transplantation date through the last date of available data;

Main study objective:

The primary objective of this study was to evaluate the FEV1 trajectory of patients diagnosed with CLAD BOS.

Study Design

Non-interventional study design

Other

Non-interventional study design, other

Retrospective chart review study

Population studied

Short description of the study population

Adult patients aged ≥ 18 years who are recipients of a lung transplant and diagnosed with CLAD-BOS following transplantation.

Age groups

- **Adult and elderly population (≥ 18 years)**
 - Adults (18 to < 65 years)
 - Adults (18 to < 46 years)
 - Adults (46 to < 65 years)
 - Elderly (≥ 65 years)
 - Adults (65 to < 75 years)
 - Adults (75 to < 85 years)
 - Adults (85 years and over)
-

Estimated number of subjects

284

Study design details

Setting

The study was conducted at approximately 6 sites located in the US and EU regions, with patient enrollment distributed nearly equally between the 2 regions. Eligible patients were identified through retrospective chart reviews of medical records at the participating study sites. The records were screened chronologically, starting with patients diagnosed with CLAD BOS from 01 January 2013. Patients who met the eligibility criteria were included in the study.

Comparators

Not applicable

Outcomes

Primary Outcome: To evaluate the FEV1 trajectory of patients diagnosed with CLAD-BOS; Secondary Outcomes: To describe demographics of patients diagnosed with CLAD-BOS, Lung Transplant History, Best Post-transplant Spirometry Parameters, CLAD-BOS Characteristics, Clinical Characteristics Following Lung Transplantation, Signs, Symptoms, or Other Clinical Findings, Trajectory of Other Relevant Spirometry Parameters, Overall Survival, Time to First CLAD-BOS Progression, Cumulative Incidence of CLAD-BOS Progression, Rate of Hospitalization due to Respiratory Failure After CLAD-BOS Onset, Incidence of Concomitant Respiratory and Non-Respiratory Disease, Use of Concomitant Treatments and Procedures for CLAD-BOS.

Data analysis plan

Descriptive summary statistics for continuous variables will include the number of patients, mean, standard deviation, median, interquartile range, and range. Descriptive summary statistics for categorical variables will include frequency counts and percentages. Confidence intervals around mean or proportion will be presented where applicable. The trajectories of FEV1 and other longitudinal

parameters will be characterized using a Linear Mixed Model for repeated measurements, using all observed available FEV1 measurements up to the last date of data available.

Overall survival and time to first CLAD-BOS progression will be assessed using Kaplan-Meier methods. Rates at yearly intervals will be reported along with the estimated medians and the corresponding CIs, whenever estimable. Kaplan-Meier curves will be presented along with the number of patients at risk at exact time points. Incidence rates will be evaluated using negative binomial or Poisson (in case of lack of convergence) regression, including an offset for patient follow-up time. Estimations will be provided along with their 95% CIs. Subgroup analyses of main outcomes will be performed by age class, gender, race, region, and severity of disease. Statistical details and potential new subgroup definitions will be reported in the Statistical Analysis Plan (SAP). The SAP will be issued before starting the collection of data with more technical and detailed elaboration of the principal features of statistical analyses. Additional post-hoc analyses may be produced to further explore data according to the results obtained. Any deviations from the original SAP (including unplanned analyses) will be documented in the Clinical Study Report.

Summary results

This study provides the most comprehensive description to date of the natural history and progression of CLAD-BOS following lung transplantation. It highlights a significant decline in lung function among patients with CLAD-BOS over a 3-year period, with spirometry parameters such as FEV₁, FVC, FEV₁/FVC, and FEF₂₅₋₇₅% showing progressive deterioration. The analysis confirms that, after disease onset, patients experience a rapid initial decline in lung function during the first 24 weeks, followed by a slower, progressive deterioration over subsequent years. This pattern was consistent across analytical approaches. Sensitivity analyses further supported the robustness of these findings. Post-hoc analyses using the clinically diagnosed date of CLAD-BOS as the starting

point, instead of the pre-specified algorithmic onset date, confirmed the same disease trajectory, an initial steep decline followed by slower progression, with only minor differences in estimated values. This consistency reinforces the reliability of the overall disease course characterization. Slight variations were observed, including a marginally greater FEV₁ decline at early timepoints and a shorter median OS (approx. 10 months less), when analyses were anchored to the clinical diagnosis date. Survival analyses demonstrated a clear difference by transplant type, with markedly longer median OS in bilateral lung transplant recipients compared to single lung recipients. The overall median survival from CLAD-BOS onset was 77.5 months, consistent with previously reported outcomes in this population. In summary, the study delineates a consistent pattern of disease progression characterized by an early, steep decline in lung function followed by stabilization or gradual deterioration over time. These data clearly indicate that the optimal therapeutic window for treating CLAD-BOS is at the early stage of disease diagnosis and provide a strong scientific rationale for in these patients as early as possible.

Data management

ENCePP Seal

The use of the ENCePP Seal has been discontinued since February 2025. The ENCePP Seal fields are retained in the display mode for transparency but are no longer maintained.

Data sources

Data sources (types)

Electronic healthcare records (EHR)

Use of a Common Data Model (CDM)

CDM mapping

No

Data quality specifications

Check conformance

Unknown

Check completeness

Unknown

Check stability

Unknown

Check logical consistency

Unknown

Data characterisation

Data characterisation conducted

Not applicable