

1. Title Page

Title	Antipsychotics in pregnancy and the risk of adverse pregnancy outcomes - a nationwide study
Research question & Objectives	1) To assess the association of antipsychotic medication use with miscarriages (primary) 2) To assess the association of antipsychotic medication use with other adverse pregnancy outcomes (secondary)
Protocol version	V2
Last update date	28/02/2025
Contributors	Primary investigator: Prof Hedvig Marie Egeland Nordeng, PhD. Professor and Head, PharmacoEpidemiology and Drug Safety (PharmaSafe) Research Group, Department of Pharmacy, University of Oslo, Oslo, Norway Contributor names: Marleen van Gelder, PhD. PharmacoEpidemiology and Drug Safety (PharmaSafe) Research Group, Department of Pharmacy, University of Oslo, Oslo, Norway Takamasa Sakai, PhD. Assistant Professor, Drug Informatics, Faculty of Pharmacy, Meijo University, Nagoya, Japan Visiting Researcher, University of Oslo, Oslo, Norway
Study registration	Site: We registered the study in the HMA-EMA Catalogue of real-world data studies. Identifier: EUPAS1000000134
Sponsor	Organization: iAPOGEE program, Pharmacoepidemiology and Drug Safety Research group, Department of Pharmacy, University of Oslo, Norway Contact: Hedvig Nordeng
Conflict of interest	None declared

Table of contents

1. Title Page	1
2. Abstract.....	3
3. Amendments and updates	4
4. Milestones	5
Table 1. Milestones.....	5
5. Rationale and background	6
6. Research question and objectives	7
Table 2. Primary and secondary research questions and objective	7
7. Research methods	9
7.1. Study design.....	9
7.2. Study design diagram.....	10
7.3. Setting.....	12
7.3.1 Context and rationale for definition of time 0 (and other primary time anchors) for entry to the study population.....	12
Table 3. Operational Definition of Time 0 (index date) and other primary time anchors	12
7.3.2 Context and rationale for study inclusion criteria:.....	13
Table 4. Operational Definitions of Inclusion Criteria	13
7.4. Variables.....	19
7.4.1 Context and rationale for exposure(s) of interest.....	19
Table 5. Operational Definitions of Exposure	19
7.4.2 Context and rationale for outcome(s) of interest	21
Table 6. Operational Definitions of Outcome.....	21
7.4.3 Context and rationale for covariates (confounding variables and effect modifiers, e.g. risk factors, comorbidities, comedications).	23
Table 7. Operational Definitions of Covariates	35
7.5. Data analysis	40
7.5.1 Context and rationale for analysis plan.....	40
Table 8. Primary, secondary, and subgroup analysis specification.....	40
Table 9. Sensitivity analyses – rationale, strengths and limitations.....	41
7.6. Data sources.....	42
7.6.1 Context and rationale for data sources.....	44
Table 10. Metadata about data sources and software.....	44
7.7. Data management	45
7.8. Quality control	45
7.9. Study size and feasibility	46
8. Limitation of the methods	49
9. Protection of human subjects.....	49
10. Reporting of adverse events	49
11. References.....	49
12. Appendices.....	52

2. Abstract

Maternal use of antipsychotics is increasing in recent years, and occurs in 0.28 to 4.64 % of all pregnancies¹⁾. Prior studies on antipsychotic safety during pregnancy have mainly focused on congenital malformations, indicating no increased risks^{2,3)}. However, less is known about other pregnancy outcomes, including miscarriages. In a prior Denmark registry study, it was reported that women who used antipsychotic medications during pregnancy had a higher risk of miscarriages compared to unexposed women (adjusted relative risk (aRR) [95% confidence interval(95%CI)]: 1.34[1.22-1.46]), but a similar risk compared to women exposed prior to (but not during) pregnancy (1.04[0.93-1.17]). Higher antipsychotic dosage (>50% of DDD) had a higher risk of miscarriages compared to unexposed women (3.19 [2.65-3.84])⁴⁾. Questions remain as to the risk of miscarriages among women who use antipsychotics in early pregnancy, also due to the methodological challenges of studying miscarriages as an outcome. Therefore, using a novel pregnancy algorithm that captures early non-live births, we aim to assess the association of second-generation antipsychotic use during pregnancy with miscarriages. In addition, we will assess associations with the other maternal and pregnancy outcomes.

We will use Norwegian nationwide registry data, which consist of the Medical Birth Registry of Norway (MBRN), linked to the Norwegian Prescription Database (NorPD) covering all dispensed medications to outpatients, the Norwegian control and payment of health reimbursements (KUHR) covering primary care contacts and the Norwegian Patient Registry (NPR) covering secondary care contacts through the maternal personal identification number. Identification of pregnancy episodes and outcomes will be done using the pregnancy algorithm developed by PharmaSafe research group at UiO⁵⁾. In brief, all pregnancies lasting ≥ 12 weeks will be identified in the MBRN, whereas primary and secondary care registries will identify pregnancies lasting < 12 weeks.

The primary exposure group is defined as second-generation antipsychotics during early pregnancy (23 days prior to gestational week 20 / end of pregnancy). Several comparison groups will be employed: 1. Unexposed, diseased comparison group, 2. First-generation antipsychotics during pregnancy (Active comparator), 3. Exposed to second-generation antipsychotics only prior to pregnancy (Discontinuer). The primary outcome is defined as miscarriages and elective termination is considered a competing outcome. We will estimate the hazard ratio with 95% CI with each comparator group, while controlling for measured confounders identified using Directed Acyclic Graphs.

In the secondary analysis, we will restrict to pregnancies identified in the MBRN. We will assess the secondary outcomes: preterm birth, small-for-gestational age (SGA), low Apgar score, transfer to NICU, congenital malformations, caesarean section, gestational diabetes, and preeclampsia.

3. Amendments and updates

Version date	Version number	Section of protocol	Amendment or update	Reason
30/04/2024	1	Final first draft		
28/02/2024	2	First revision	Updated mainly on covariates information and inclusion/exclusion criteria	Reflected the latest findings on the missing value imputations in medical birth registries Refinement of eligibility criteria

4. Milestones

Table 1. Milestones

Milestone	Planned date
Feasibility counts	30/09/2023
Protocol finalized	30/04/2024
Registration of protocol	30/04/2024
Final report of study results	28/02/2026

5. Rationale and background

What is known about the condition: Psychiatric conditions for which antipsychotics are prescribed include schizophrenia, bipolar disorders and depressive disorders. These conditions have all been associated with adverse pregnancy outcomes⁶⁻⁹. For example, in a French population-based study, women with schizophrenia experienced more pregnancy complications (aOR[95%CI]: 1.41[1.31–1.51]), delivery complications (1.18[1.09–1.29]), and caesarean sections (1.15[1.05–1.25]) compared with women without severe mental disorders¹⁰. In the same study, newborns of women with schizophrenia had more neonatal complications (1.38[1.27–1.50]), were more commonly born preterm (1.64[1.42–1.90]), small-for-gestational-age (1.34[1.19–1.50]) and with a low birth weight (1.75[1.53–2.00])¹⁰. In an Australian population-based cohort study, women with severe mental illness had higher risks of gestational diabetes (aOR[95%CI]:1.57 [1.34–1.84]), unplanned caesarean section (1.17 [1.02–1.33]), having a newborn with a low Apgar score at 5 minutes (1.50 [1.19–1.90]), preterm birth (1.40 [1.20–1.63]), and low birth weight (1.26 [1.06–1.49])¹¹. Furthermore, in a Norwegian registry study, some psychiatric disorders have been associated with an increased risk of miscarriages after adjustment for co-occurring psychiatric disorders with an aOR of 1.22 [95%CI 1.03–1.44] for schizophrenia spectrum disorders, 1.27 [1.19–1.36] for bipolar disorders, and 1.21 [1.19–1.23] for depressive disorders). This risk was further increased among women with more than one psychiatric diagnosis (aOR[95%CI]: 1.45 [1.40–1.51] for two psychiatric diagnoses, 1.51 [1.31–1.73] for three or more diagnoses)¹².

What is known about the exposure of interest: In many countries, the prevalence of antipsychotic use during pregnancy is increasing, especially for second-generation antipsychotics, ranging typically between 0.17 and 1.53%¹. In Norway (2005–2015), the prevalence of antipsychotic use during the pregnancy period was 1.16% (second-generation antipsychotics: 0.24%, first-generation antipsychotics: 0.95%)¹.

From Denmark, it is reported that women who used antipsychotic medication during pregnancy had a 34% higher risk of miscarriages (aRR [95%CI]: 1.34[1.22; 1.46]) compared to unexposed women, but a similar risk compared to women exposed prior to (but not during) pregnancy (1.04[0.93; 1.17]). Higher antipsychotic dosage (>50% of DDD) had a higher risk of miscarriages compared to unexposed women (3.19 [2.65; 3.84])⁴. From the analysis of a Japanese spontaneous reporting database, a potential signal for miscarriages was detected for aripiprazole [reporting OR [95%CI]: 2.76 [1.62–4.69]; n = 18]. In contrast, no potential signal for miscarriages was detected for other second-generation antipsychotics¹³. Post-marketing surveillance studies on olanzapine, quetiapine, and risperidone showed no associations with miscarriages^{14–16}.

Gaps in knowledge: Questions remain as to the risk of miscarriages among women who use antipsychotics in early pregnancy. More specifically, the risks of different antipsychotics classes (first-generation antipsychotics, second-generation antipsychotics) are unknown.

What is the expected contribution of this study? This study using nation-wide registries from Norway contributes to the safety information of second-generation antipsychotic use during pregnancy on birth outcomes other than congenital malformations.

6. Research question and objectives

Table 2. Primary and secondary research questions and objective

A. Primary research question and objective

Objective:	To evaluate the association of exposure to second-generation antipsychotics during pregnancy with the risk of miscarriages while taking into account competing risks of elective terminations.
Hypothesis:	The risk of miscarriages is elevated with second-generation antipsychotic use during pregnancy.
Population (<i>mention key inclusion-exclusion criteria</i>):	<p>All pregnancies identified in the MBRN (Medical Birth Registry of Norway) for pregnancies lasting ≥ 12 weeks, and primary and secondary care registries for pregnancies lasting < 12 weeks.</p> <p>We will use the recently developed pregnancy algorithm by PharmaSafe researchers at UiO including approximately 860,000 pregnancies (2008-2018), including live-births (74.8%), miscarriages (13.1%), elective terminations (11.0%), ectopic pregnancies (0.7%), stillbirths (0.3%), and molar pregnancies (0.1%). Ectopic and molar pregnancies will be excluded.</p> <p>Of these pregnancies, we restrict mothers having a mental disease diagnostic code or antipsychotic dispensing between 6 months prior to pregnancy and gestational week 4 (index date).</p>
Exposure:	Second-generation antipsychotics during early pregnancy (23 days prior to gestational week 20 / end of pregnancy)
Comparator:	<ol style="list-style-type: none"> 1. Unexposed, diseased-comparison group (Antipsychotics unexposed before and during early pregnancy) 2. Active comparator (First-generation antipsychotics during early pregnancy) 3. Discontinuer (Exposed to second-generation antipsychotics only prior to pregnancy not during early pregnancy)
Outcome:	Miscarriages
Time (<i>when follow up begins and ends</i>):	From the 4 weeks of gestation to 20 weeks of gestation
Setting:	Primary and secondary care (Dx), Outpatient (Rx), MBRN information
Main measure of effect:	Hazard Ratio with 95% confidence intervals

B. Secondary research question and objective

Objective:	To evaluate the association of exposure to second-generation antipsychotics during pregnancy with the risk of selected pregnancy outcomes other than miscarriages
Hypothesis:	The risk of other selected pregnancy outcomes is elevated with second-generation antipsychotic use during pregnancy.
Neonatal outcomes	
Population (<i>mention key inclusion-exclusion criteria</i>):	All pregnancies identified in the MBRN (Medical Birth Registry of Norway) Of these pregnancies, we restrict mothers having a mental disease diagnostic code or antipsychotic dispensing up to 6 months prior to pregnancy.
Exposure:	Second-generation antipsychotics during pregnancy (congenital malformations: during first trimester, preterm birth: until gestational week 37)
Comparator:	<ol style="list-style-type: none"> 1. Unexposed, diseased-comparison group (Antipsychotics unexposed before and during pregnancy) 2. Active comparator (First-generation antipsychotics during pregnancy) 3. Discontinuer (Exposed to second-generation antipsychotics only prior to pregnancy not during pregnancy)
Outcome:	Preterm birth, small-for-gestational-age (SGA), low Apgar score, transfer to NICU, congenital malformations
Time (<i>when follow up begins and ends</i>):	<p>From the start of the pregnancy to the delivery date: preterm birth, SGA, low Apgar score</p> <p>From the start of the pregnancy to the discharge: transfer to NICU</p> <p>From the start of the pregnancy to 1 year after birth: congenital malformations</p>
Setting:	Primary and secondary care (Dx), Outpatient (Rx), MBRN information
Main measure of effect:	Relative Risk with 95% confidence intervals
Maternal outcomes	
Population (<i>mention key inclusion-exclusion criteria</i>):	All pregnancies identified in the MBRN (Medical Birth Registry of Norway) Of these pregnancies, we restrict mothers having a mental disease diagnostic code or antipsychotic dispensing up to 6 months prior to pregnancy.
Exposure:	Second-generation antipsychotics during pregnancy
Comparator:	<ol style="list-style-type: none"> 1. Unexposed, diseased-comparison group (Antipsychotics unexposed before and during pregnancy)

	2. Active comparator (First-generation antipsychotics during pregnancy) 3. Discontinuer (Exposed to second-generation antipsychotics only prior to pregnancy)
Outcome:	Gestational diabetes, preeclampsia, caesarean section
Time (<i>when follow up begins and ends</i>):	From the start of the pregnancy to the delivery date: gestational diabetes, preeclampsia, caesarean section,
Setting:	Primary and secondary care (Dx), Outpatient (Rx), MBRN information
Main measure of effect:	Relative Risk with 95% confidence intervals

7. Research methods


7.1. Study design

Research design: Cohort study


Rationale for study design choice: Since we are planning to study a single exposure in relation to multiple outcomes, we will use a cohort design.

7.2. Study design diagram

Primary outcome



Secondary outcomes



7.3. Setting

7.3.1 Context and rationale for definition of time 0 (and other primary time anchors) for entry to the study population

In this study, we will use the date of 4 weeks of gestation as time 0 for the primary outcome. This is to exclude unnoticeable pregnancies and to avoid misclassification of potential discontinuers just after recognizing being pregnant.

For the secondary outcome, we will use the date of start of the pregnancy (last menstrual period, LMP) as time 0.

For the primary analysis, follow-up ends at either end of pregnancy or 20 weeks of gestation, whichever comes first.

Table 3. Operational Definition of Time 0 (index date) and other primary time anchors

Study population name(s)	Time Anchor Description (e.g. time 0)	Number of entries	Type of entry	Washout window	Care Setting ¹	Code Type ²	Diagnosis position	Incident with respect to...	Measurement characteristics/ validation	Source of algorithm
Pregnant women	Start of the pregnancy	Multiple (women with multiple pregnancies are allowed to contribute multiple times)	Incident	See pregnancy algorithm	See pregnancy algorithm	ICD-10, ICPC-2, MBRN variables	See pregnancy algorithm	-		MBRN, NPR, KUHR

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

²See appendix for listing of clinical codes for each study parameter

7.3.2 Context and rationale for study inclusion criteria:

Pregnancies identified with the UiO pregnancy algorithm using MBRN for pregnancies lasting ≥ 12 weeks and primary and secondary care registries for pregnancies lasting < 12 weeks will be included. To ensure that there is sufficient observable time, women must have registry coverage from 6 months before their last menstrual period to end of pregnancy.

Table 4. Operational Definitions of Inclusion Criteria

Criterion	Details	Order of application	Assessment window	Care Settings ¹	Code Type ²	Diagnosis position ³	Applied to study populations:	Measurement characteristics/validation	Source for algorithm
Primary outcome									
Pregnancies identified in MBRN, NPR, KUHR		Before selection of study population	See pregnancy algorithm	See pregnancy algorithm	See pregnancy algorithm	See pregnancy algorithm	Pregnant women		pregnancy algorithm ⁵⁾
Observable time	Mothers with observable time from 6 month before LMP (NPR registers) to 10 months after LMP	Before selection of study population	Mother: [LMP-180, LMP+10 months]	n/a	n/a	n/a	Pregnant women	n/a	n/a
Psychiatric condition between 6 months prior to pregnancy and gestational week 4	Psychiatric disorder diagnosis and/or antipsychotic dispensing between 6 months prior to pregnancy and gestational week 4	After selection of study population by other eligibility criteria	[LMP-180, GW4]	All	ATC code ICD-10 code/ICPC -2 code	n/a	Pregnant women	n/a	Investigator defined
Secondary outcomes									
Pregnancies identified in MBRN (Congenital malformation, Maternal outcomes other than caesarean section)	Identified from MBRN	After selection of study population by other eligibility criteria	n/a	All	n/a	n/a	Pregnant women		MBRN ¹⁷⁾
Singleton live birth identified in MBRN (Caesarean section, Neonatal outcomes)	Identified from MBRN	After selection of study population by other eligibility criteria	[EoP, EoP]	All	n/a	n/a	Pregnant women		MBRN ¹⁷⁾

other than congenital malformation)									
Observable time (Congenital malformations)	Mother from 6 month before LMP (NPR registers) Child to 1 year after delivery	Before selection of study population	Mother: [LMP-180, EoP] Child: [EoP, EoP+1y]	All	n/a	n/a	Pregnant women	n/a	n/a
Psychiatric condition 6 months prior to pregnancy	Psychiatric disorder diagnosis and/or antipsychotic dispensing 6 months prior to pregnancy	After selection of study population by other eligibility criteria	[LMP-180, LMP]	All	ATC code ICD-10 code/ICPC-2 code	n/a	Pregnant women	No validation study	Investigator defined

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

7.3.1.1. Appendix A. Codes for the psychiatric disorder

Covariates		Data source	Code (ICD-10 code/ICPC-2 code)
Psychiatric disorder	Schizophrenia	NPR/KUHR	ICD-10 code: F20-F29 ICPC-2 code: P72, P73, P98
	Bipolar disorder		ICD-10 code: F31
	Mania		ICD-10 code: F30
	Depressive disorder with psychotic symptoms		ICD-10 code: F323, F333

7.3.1.2. Appendix B. ATC codes for the drug exposure

Group of antipsychotics (Investigator defined)		Data source	ATC code
Second-generation antipsychotics	Sertindole	NorPD	N05AE03
	Ziprasidone		N05AE04
	Lurasidone		N05AE05
	Clozapine		N05AH02
	Olanzapine		N05AH03
	Quetiapine		N05AH04
	Asenapine		N05AH05
	Sulpiride		N05AL01
	Sultopride		N05AL02

	Remoxipride		N05AL04
	Amisulpride		N05AL05
	Levosulpiride		N05AL07
	Risperidone		N05AX08
	Mosapramine		N05AX10
	Zotepine		N05AX11
	Aripiprazole		N05AX12
	Paliperidone		N05AX13
	Iloperidone		N05AX14
	Cariprazine		N05AX15
	Brexpiprazole		N05AX16
First-generation antipsychotics	Chlorpromazine		N05AA01
	Levomepromazine		N05AA02
	Promazine		N05AA03
	Acepromazine		N05AA04
	Triflupromazine		N05AA05
	Cyamemazine		N05AA06
	Chlorproethazine		N05AA07
	Dixyrazine		N05AB01
	Fluphenazine		N05AB02
	Perphenazine		N05AB03
	Prochlorperazine		N05AB04
	Thiopropazate		N05AB05
	Trifluoperazine		N05AB06
	Acetophenazine		N05AB07
	Thiopropazine		N05AB08
	Butaperazine		N05AB09
	Perazine		N05AB10
	Periciazine		N05AC01
	Thioridazine		N05AC02
	Mesoridazine		N05AC03
	Pipotiazine		N05AC04
	Haloperidol		N05AD01
	Trifluperidol		N05AD02
	Melperone		N05AD03
	Moperone		N05AD04
	Pipamperone		N05AD05
	Bromperidol		N05AD06
	Benperidol		N05AD07
	Droperidol		N05AD08
	Fluanisone		N05AD09
	Oxypertine		N05AE01
	Molindone		N05AE02
	Flupentixol		N05AF01
	Clopentixol		N05AF02
	Chlorprothixene		N05AF03

	Tiotixene		N05AF04
	Zuclopenthixol		N05AF05
	Fluspirilene		N05AG01
	Pimozide		N05AG02
	Penfluridol		N05AG03
	Loxapine		N05AH01
	Clotiapine		N05AH06
	Tiapride		N05AL03
	Veralipride		N05AL06
	Prothipendyl		N05AX07

***Note**

The ATC codes are cited from previous drug utilization study:

Reutfors J, Cesta CE, Cohen JM, et al. Antipsychotic drug use in pregnancy: A multinational study from ten countries. Schizophr Res. 2020 Jun;220:106-115. PMID: 32295750.

7.3.3 Context and rationale for study exclusion criteria

Pregnancies with no possibility of livebirth (e.g. ectopic and molar pregnancies) will be excluded from the primary analysis.

Table 5. Operational Definitions of Exclusion Criteria

Primary outcome

Criterion	Details	Order of application	Assessment window	Care Settings ¹	Code Type ²	Diagnosis position ³	Applied to study populations:	Measurement characteristics/ validation	Source for algorithm
Ectopic and molar pregnancies	NPR for identification	After selection of study population by inclusion criteria	[LMP, GW12]		ICD-10		Pregnant women	Pregnancy algorithm ⁵⁾	
Implausible early dating pregnancies	Pregnancies less than 4 weeks	After selection of study population by eligibility criteria	[LMP, GW4]				Pregnant women	Pregnancy algorithm ⁵⁾	

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

Secondary outcomes

Criterion	Details	Order of application	Assessment window	Care Settings ¹	Code Type ²	Diagnosis position ³	Applied to study populations:	Measurement characteristics/ validation	Source for algorithm
known teratogen exposure (Neonatal outcomes)		After selection of study population by inclusion criteria	[LMP, EoP-1]		ATC code		Pregnant women		n/a
pre-existing diabetes (Gestational diabetes)	MBRN: DIABETES_MELLITUS Definition: 1: type 1 diabetes diagnosed prior to pregnancy 2: type 2 diabetes diagnosed prior to pregnancy 3: other or unspecified diabetes diagnosed prior to pregnancy	After selection of study population by inclusion criteria		IP, OP	Using variable in MBRN		Pregnant women		MBRN ¹⁷⁾

7.3.1.3. Appendix C. List of teratogenic medication

Teratogens	ATC code	Data source
Antiepileptics		NorPD
Carbamazepine	N03AF01	
Phenobarbital	N03AA02	
Primidone	N03AA03	
Valproic acid	N03AG01	
Phenytoin	N03AB02	
	N03AB52	
Fosphenytoin	N03AB05	
Topiramate	N03AX11	
Retinoids		
Acitretin	D05BB02	
Alitretinoin	D11AH04	
Bexarotene	L01XF03	
Isotretinoin	D10BA01	
Tretinoin	L01XF01	
Etretinate	D05BB01	
Anti-thyroids		
Carbimazole	H03BB01	
Thiamazole	H03BB02	
Normothymic		
Lithium	N05AN01	
Anticoagulants		
Warfarin	B01AA03	
Phenindione	B01AA02	
Acenocumarol	B01AA07	
Immunomodulants		
Mycophenolate	L04AA06	
Fingolimod	L04AA27	
Pomalidomide	L04AX06	
Lenalidomide	L04AX04	
Cytotoxic		
Cisplatin	L01XA01	
Doxorubicin	L01DB01	
Methotrexate	L04AX03	
	L01BA01	
Thalidomide	L04AX02	
Hormonal		
Diethylstilbestrol	L02AA01	
	G03CB02	
	G03CC05	
Misoprostol	A02BB01	
	G02AD06	

7.4. Variables

7.4.1 Context and rationale for exposure(s) of interest

The lack of randomization has to be compensated for by methods that maximize the comparability between the exposed and comparison groups. Unexposed to any antipsychotics with psychiatric disorder (listed in Appendix A) is a disease-matched control group. First-generation antipsychotics are used as active comparators for second-generation antipsychotics which have the overlap of some indication (schizophrenia, mania and bipolar disorder). Past users during 180 days before the pregnancy but unexposed during pregnancy is defined as discontinuers. The selection of the comparison group will not be the only method to increase exchangeability between the exposed and unexposed groups; we will also adjust for measured confounders.

Table 6. Operational Definitions of Exposure

Primary Outcome

The lag time between arrest of development and miscarriages (median 23 days)¹⁸⁾ is considered in the assessment windows.

Exposure group name(s)	Detail	Washout window	Assessment Window	Case Setting ¹	Code Type ²	Dagnosis position ³	Applied to study populations	Incident with respect to...	Measurement characteristics/validation	Source of algorithm
Exposure: Second-generation antipsychotics	Second-generation antipsychotics exposure as listed in Appendix B		[GW4, EoP/20w-23]**	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined
Comparator 1: Unexposed, diseased-comparison group	No antipsychotic up to 6 months prior to pregnancy and during pregnancy.	[LMP-180, LMP]	[LMP-180, EoP/20w-23]**	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined
Comparator 2: Active comparator	First-generation antipsychotics exposure as listed in Appendix B (Active comparator)		[GW4, EoP/20w-23]**	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined
Comparator 3: Discontinuer	Non-use of antipsychotics after GW4 but past users during 6 months before the start of the pregnancy or before GW4 in pregnancy(Discontinuer)		[LMP-180, EoP/20w-23]**	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

* Definition of psychiatric disorder will be listed in appendix B

** Earlier date either 23 days prior to end of pregnancy or 20 weeks of gestation.

Secondary Outcome

Exposure group name(s)	Detail	Washout window	Assessment Window [†]	Care Setting ¹	Code Type ²	Diagnosis position ³	Applied to study populations	Incident with respect to...	Measurement characteristics/validation	Source of algorithm
Exposure: Second-generation antipsychotics	Second-generation antipsychotics exposure as listed in Appendix C		[LMP, EoP-1]	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined
Comparator 1: Unexposed, diseased-comparison group	No antipsychotic up to 6 months prior to pregnancy and during pregnancy.	[LMP-180, LMP]	[LMP-180, EoP-1]	n/a	ATC code ICPC-2 code ICD-10	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined
Comparator 2: Active comparator	First-generation antipsychotics exposure as listed in Appendix C (Active comparator)		[LMP, EoP-1]	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined
Comparator 3: Discontinuer	Non-use of antipsychotics after LMP but past users during 6 months before the start of the pregnancy (Discontinuer)		[LMP-180, EoP-1]	n/a	ATC code	n/a	Pregnant women with psychiatric disorder*		No validation study	Investigator defined

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

* Definition of psychiatric disorder will be listed in Appendix B

[†] Exposure assessment window: congenital malformations: during first trimester, preterm birth: until 37 week

Appendix B. Codes for the psychiatric disorder

Appendix C. ATC codes for the drug exposure

7.4.2 Context and rationale for outcome(s) of interest

Little is known about the risk of miscarriages following the use of second-generation antipsychotics in early pregnancy.

Table 7. Operational Definitions of Outcome

Primary analysis

Outcome name	Details	Primary outcome?	Type of outcome	Washout window	Care Settings ¹	Code Type ²	Diagnosis Position ³	Applied to study populations:	Outcome measurement characteristics/ validation	Source of algorithm
Miscarriages	Days from cohort entry date, numerical	Yes	time-varying (binary)	n/a	IP, OP	Using variables in MBRN, NPR, KUHR	Primary	Pregnant women		Pregnancy algorithm ⁵⁾
Elective termination	As a competing outcome of miscarriages Days from cohort entry date, numerical	No	time-varying (binary)	n/a	IP, OP	Using variables in MBRN, NPR	Primary	Pregnant women		Pregnancy algorithm ⁵⁾

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

Secondary analysis

Outcome name	Details	Primary outcome?	Type of outcome	Washout window	Care Settings ¹	Code Type ²	Diagnosis Position ³	Applied to study populations :	Outcome measurement characteristics/ validation	Source of algorithm
Preterm birth	Length of gestation in whole weeks, numerical (SVLEN) <37	Yes	binary	n/a	IP, OP	Using variables in MBRN		Neonate	Acta Obstet Gynecol Scand. 2016;95(5):519-527.	MBRN ¹⁷⁾
Small-for-gestational-age	Defined as the 10% of children with the lowest birth weight given gestational age. Deviations from the expected birth weight for the child given gestational age and child sex,	Yes	binary	n/a	IP, OP	Using variables in MBRN		Neonate		MBRN ¹⁷⁾

	numerical (ZSCORE_BW_GA) <- 1.28 (lower 10 percentile value)									
Low Apgar score	Apgar score at 5 minutes after birth (APGAR5) < 7	Yes	binary	n/a	IP, OP	Using variables in MBRN		Neonate		MBRN ¹⁷⁾
Transfer to NICU	Child transferred to neonatal intensive care unit (OVERFLYTET) Definition: 1: yes	Yes	binary	n/a	IP, OP	Using variables in MBRN		Neonate		MBRN ¹⁷⁾
Congenital malformations	Child congenital malformation, any (MISD) Definition: 1: yes	Yes	binary	n/a	IP, OP	Using variables in MBRN		Infant		MBRN ¹⁷⁾ EUROCAT Guide 1.4: Instruction for the registration of congenital anomalies: EUROCAT Central registry, University of Ulster, 2013
Caesarean section	we will utilize following variable to code the unplanned caesarean section binary; Caesarean section (KSNITT) Definition: KSNITT != Null 1: planned caesarean 2: emergency caesarean 9: unspecified caesarean	Yes	binary	n/a	IP, OP	Using variables in MBRN		Pregnant women	Acta Obstet Gynecol Scand. 2017;96(7):892-897.	MBRN ¹⁷⁾
Gestational diabetes	Maternal diabetes (DIABETES_MELLITUS) Definition: 4: gestational diabetes	Yes	binary	n/a	IP, OP	Using variables in MBRN		Pregnant women		MBRN ¹⁷⁾
Preeclampsia	Early onset preeclampsia (PREEKLIDL) 1: yes Preeclampsia – a hypertensive disorder of pregnancy (PREEKL) 1: mild, 2: severe, 3: unspecified Definition: PREEKL != Null or PREEKLIDL == 1	Yes	binary	n/a	IP, OP	Using variables in MBRN		Pregnant women	Acta Obstet Gynecol Scand 2013;92(8):943-950.	MBRN ¹⁷⁾

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

7.4.3 Context and rationale for follow up

Table 8. Operational Definitions of Follow Up

A. Primary analysis

Follow up start	Start of the pregnancy (LMP)	
Follow up end¹	Select all that apply	Specify
Date of outcome	Yes	Including competing outcomes
Date of death	No	
End of observation in data	No	
Day X following index date (specify day)	Yes	20 weeks of gestation
End of study period (specify date)	No	
End of exposure (specify operational details, e.g. stockpiling algorithm, grace period)	No	
Date of add to/switch from exposure (specify algorithm)	No	
Other date (specify)	No	

¹ Follow up ends at the first occurrence of any of the selected criteria that end follow up.


B. Secondary analysis

Follow up start	Start of the pregnancy (LMP)	
Follow up end ¹	Select all that apply	Specify
Date of outcome	Yes	
Date of death	No	
End of observation in data	No	
Day X following index date (specify day)	Yes	transfer to NICU: until the discharge congenital malformations: 1 year after birth
End of study period (specify date)	No	
End of exposure (specify operational details, e.g. stockpiling algorithm, grace period)	No	
Date of add to/switch from exposure (specify algorithm)	No	
Other date (specify)	No	

7.4.4 Context and rationale for covariates (confounding variables and effect modifiers, e.g. risk factors, comorbidities, comedications).


We will measure maternal characteristics including baseline sociodemographic characteristics, maternal comorbidities, and concomitant medications, and pregnancy characteristics. These were selected due to their potential to act as confounders or effect modifiers based on following the directed acyclic graph (DAG) below. In order to avoid including potential intermediate variables into covariates, we basically included information before the index date.

Primary outcome:
Miscarriages




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Secondary outcomes:
Preterm birth




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Small-for-gestational-age




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Low Apgar score




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Transfer to NICU




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Congenital malformations




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Gestational diabetes




We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Preeclampsia



We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Caesaren section



We will use the psychiatric disorder-related outpatient visit information as a proxy for psychiatric disorder severity.

Table 9. Operational Definitions of Covariates

Characteristic	Details	Type of variable	Assessment window	Care Settings ¹	Code Type ²	Diagnosis Position ³	Applied to study populations:	Measurement characteristics/validation	Source for algorithm
Maternal characteristics									
Maternal age	MORS_ALDER	Continuous	[LMP, LMP]	n/a	MBRN (numerical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Marital status	SIVST Married/cohabiting Dichotomize(Yes/No) As socioeconomic background	Binary	[LMP, LMP]	n/a	MBRN (categorical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Maternal employment status	YRKE_KODE Dichotomize(Yes/No) 1: not employed 2: full time employed 3: part time employed As socioeconomic background	Binary	[LMP, LMP]	n/a	MBRN (categorical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Smoking status	ROYK_FOER (smoking before pregnancy) Dichotomize(Yes/No) 1: no 2: yes, sometimes 3: yes, daily	Binary	Primary ROYK_FOER, Secondary ROYK_FOER,	n/a	MBRN (categorical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Maternal weight	Mother's weight before pregnancy (MORS_VEKT_FOER)	Continuous	[LMP, LMP]	n/a	MBRN (numerical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Calendar year	Year of start of the pregnancy	Continuous	[LMP, LMP]	n/a	MBRN (numerical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Hypertension	Maternal hypertension diagnosed before pregnancy (HYPERTENSJON_KRONISK)	Binary	[LMP, LMP]	n/a	MBRN (categorical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Patient Register Index	Modified and updated charlson comorbidity index for NPR Refer to Appendix D	Continuous	Primary [LMP-180, GW4] Secondary	Any	NPR	n/a	Pregnant women	n/a	Clinical Epidemiology 2014:6 395–404

Characteristic	Details	Type of variable	Assessment window	Care Settings ¹	Code Type ²	Diagnosis Position ³	Applied to study populations:	Measurement characteristics/validation	Source for algorithm
			[LMP-180, LMP]						
Central Nervous System (CNS)-acting comedications	CNS-acting comedications which may be associated with miscarriages Anticonvulsants, benzodiazepines, SSRI, SNRI	Binary	Primary [LMP-180, GW4] Secondary [LMP-180, LMP]	Any	NorPD(ATC code)	n/a	Pregnant women	n/a	Investigator defined
Teratogen exposure	Refer to Appendix A	Binary	Primary [LMP-180, GW4]	Any	NorPD(ATC code)	n/a	Pregnant women	n/a	n/a
Pregnancy characteristics									
Folate use	FOLATF (Use of folate before pregnancy) Yes/No (by default)	Categorical	Primary FOLATF Secondary FOLATF	n/a	MBRN (binary)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Previous pregnancy loss	SPABORT_12_5 (Number of previous miscarriages before week 12) SPABORT_23_5 (Number of previous miscarriages/stillbirths week 12-23) DODFODTE_5 (Number of previous stillbirths after week 24) Combine the above variables, or if MBRN information is not available, we will count the number based on the UiO pregnancy algorithm. 0, 1, 2+	Categorical	[LMP, LMP]	n/a	MBRN (Categorical) And pregnancy algorithm ⁵⁾ using variables in MBRN, NPR, KUHR	n/a	Pregnant women	n/a	MBRN ¹⁷⁾ pregnancy algorithm ⁵⁾

Characteristic	Details	Type of variable	Assessment window	Care Settings ¹	Code Type ²	Diagnosis Position ³	Applied to study populations:	Measurement characteristics/validation	Source for algorithm
Previous preterm birth, gestational diabetes, preeclampsia, caesarean section	Count up the each outcomes in the previous pregnancies using the same definition of each outcomes.	Continuous	Same as the outcome section	Same as the outcome section	Same as the outcome section	Same as the outcome section	Same as the outcome section	Same as the outcome section	MBRN ¹⁷⁾
Parity	PARITET_5 (Number of previous births) 0, 1, 2, 3, 4+ Dichotomize(0/0<)	Categorical	[LMP, LMP]	n/a	MBRN (Categorical)	n/a	Pregnant women	n/a	MBRN ¹⁷⁾
Maternal infection	Refer to Appendix D	Binary	Primary [LMP-180, GW4] Secondary [LMP-180, LMP]	n/a	NPR KUHR	n/a	Pregnant women	n/a	Investigator defined
Obstetric comorbidity index	Secondary analysis only Refer to Appendix D	Continuous	[LMP, EoP]	n/a	MBRN	n/a	Pregnant women	n/a	Clin Epidemiol. 2021 Feb 26;13:161-174.
Psychiatric disorder-related characteristics									
Psychiatric disorder-related outpatient visit	As a proxy of psychiatric disorder severity	Continuous	Primary [LMP-180, GW4] Secondary [LMP-180, LMP]	OP	NPR(ICD-10) KUHR (ICD-10, ICPC-2) Refer to appendix B	Primary or secondary	Pregnant women	n/a	n/a

¹ IP = inpatient, OP = outpatient, ED = emergency department, OT = other, n/a = not applicable

² See appendix for listing of clinical codes for each study parameter

³ Specify whether a diagnosis code is required to be in the primary position (main reason for encounter)

7.4.1.1. Appendix D. Data source and codes for defining the covariates

Covariates		Data source	Code (ATC code/ICD-10 code/ICPC-2 code)
Obstetric comorbidity index		MBRN	Refer to Appendix D-1
Patient Register Index		NPR KUHR	Refer to Appendix D-2
Maternal infection		KUHR	Respiratory infection ICPC-2: R71, R72, R74, R75, R76, R77, R78, R80, R81, R83 Urinary infection ICPC-2: U70, U71, U72 Other infection ICD-10: E06.0, I30.x except I30.0, I32.0, I32.1, I33.x, I38, I39.x, I40.0, I41.0, I41.1, I41.2, I43.0, I52.0 ICPC-2: K70, L87, T70, W70, W71, W94
Teratogen exposure		NorPD	Refer to Appendix A
Central Nervous System (CNS)-acting comedications	Anticonvulsants	NorPD	N03A
	Benzodiazepines		N05BA, N05CD
	SSRI		N06AB
	SNRI		N06AX16, N06AX17, N06AX21, N06AX23, N06AX28

7.4.1.2. Appendix D-1 Obstetric Comorbidity Index

Co-morbidity	MBRN variable	Weight
Severe Pre-Eclampsia	(EKLAMPSI==1 or HELLP==1 or PREEKL==2 or PREEKLTIDL==1) & HYPERTENSJON_KRONISK!=1	5
Mild/Unspecified Pre-Eclampsia	(PREEKL==1 or PREEKL==3) & HYPERTENSJON_KRONISK!=1 & Severe Pre-Eclampsia !=1	2
Multiple Gestation	FLERFODSEL==1	2
Chronic Renal Disease	NYRESYKDOM_KRONISK==1	1
Pre-Existing Hypertension	HYPERTENSJON_KRONISK==1	1
Asthma	ASTMA==1	1
Pre-Existing Diabetes Mellitus	DIABETES_MELLITUS == 1 or 2 or 3	1
Gestational Hypertension	HYPERTENSJON_ALENE==1 & HYPERTENSJON_KRONISK!=1 & Severe Pre-Eclampsia !=1 & Mild/Unspecified Pre-Eclampsia!=1	1
Maternal age	MORS_ALDER>44	3
	40 <= MORS_ALDER <= 44	2
	35 <= MORS_ALDER <= 39	1
Previous Caesarean Delivery	Registered previous pregnancy outcomes (KSNITT)	1

7.4.1.3. Appendix D-2 Patient Register Index

Co-morbidity	ICD10 code	Weight
Myocardial infarction	I21, I22, I252	1
Congestive heart failure	I43, I50, I099, I110, I130, I255, I420, I425, I426, I427, I428, I429, P290	2
Peripheral vascular disease	I70, I71, I731, I738, I739, I771, I790, I792, K551, K558, K559, Z958, Z959	1
Cerebrovascular disease	G45, G46, I60, I61, I62, I63, I64, I65, I66, I67, I68, I69, H340	1
Dementia	F00, F01, F02, F03, G30, F051, G311	2
Chronic pulmonary disease	J40, J41, J42, J43, J44, J45, J46, J47, J60, J61, J62, J63, J64, J65, J66, J67, I278, I279, J684, J701, J703	1
Connective tissue disease	M32, M33, M34, M05, M06, M315, M351, M353, M360	1
Ulcer disease	K25, K26, K27, K28	1
Mild liver disease	B18, K73, K74, K709, K717, K760, K768, K769, Z944, K700, K701, K702, K703, K713, K715, K762, K763, K764	2
Diabetes	E100, E101, E106, E108, E109, E110, E111, E116, E118, E119, E120, E121, E126, E128, E129, E130, E131, E136, E138, E139, E140, E141, E146, E148, E149	0
Diabetes with end organ damage	E107, E117, E127, E137, E147, E102, E103, E104, E105, E112, E113, E114, E115, E122, E123, E124, E125, E132, E133, E134, E135, E142, E143, E144, E145	1
Hemiplegia	G81, G82, G041, G114, G801, G802, G839, G830, G831, G832, G833, G834	1
Moderate or severe renal disease	I120, I131, N250, Z940, Z992, N032, N033, N034, N035, N036, N037, N052, N053, N054, N055, N056, N057, Z490, Z491, Z492	1
Any tumor/leukemia/lymphoma	C0, C1, C2, C3, C5, C6, C9, C40, C41, C42, C43, C45, C46, C47, C48, C49, C70, C71, C72, C73, C74, C75, C76, C81, C82, C83, C84, C85, C88, C89	2
Moderate or severe liver disease	I850, I859, I864, I982, K704, K711, K721, K729, K765, K766, K767	5
Metastatic solid tumor	C77, C78, C79, C80	6
HIV/AIDS	B20, B21, B22, B24	3

7.5. Data analysis

7.5.1 Context and rationale for analysis plan

Table 10. Primary, secondary, and subgroup analysis specification

A. Primary analysis

Hypothesis:	Maternal use of second-generation antipsychotics during pregnancy increases the risk of miscarriage
Exposure contrast:	Second-generation antipsychotics vs Comparator 1: Unexposed, diseased-comparison group Second-generation antipsychotics vs Comparator 2: Active comparator Second-generation antipsychotics vs Comparator 3: Discontinuer
Outcome:	Miscarriage
Analytic software:	R
Model(s):	<p><u>Outcome model:</u> Cause-specific HRs¹⁹⁾ will be estimated with Cox proportional hazards regression models. We will use robust standard errors to account for correlation within women who participated with >1 pregnancy in this study. An elective termination could potentially have ended in a miscarriage, if the pregnancy had not already been terminated. This case should be censored, thereby taking competing risks into account.</p> <p><u>Propensity score model:</u> Logistic regression model, Exposure = covariates Covariates: Maternal age, Marital status, Maternal employment status, Smoking status before pregnancy, Maternal weight, Calendar year, Patient Register Index before pregnancy, CNS-acting comedications, Folate use before pregnancy, Teratogen exposure before pregnancy, Previous pregnancy loss, Parity, Maternal infection Psychiatric disorder-related outpatient visit</p>
Confounding adjustment method	Propensity scores will be estimated using logistic regression with exposure as a dependent variable and potential confounders and risk factors for the outcome as independent variables. Including all potential confounders and risk factors for the outcome in the propensity score estimation has demonstrated increased precision without increased bias. After checking the propensity score distribution, we will decide how to deal with the extreme scores. In the weighted cohort, we will estimate HRs with Cox proportional hazards regression models. Depending on the results of the exploratory data analyses, we may apply other confounding adjustment methods (e.g. pooled logistic regression) of deemed more appropriate with regard to exposure patterns and proportional hazard assumption.
Missing data methods	If information derived from the MBRN on covariates is missing, obtain the information from other pregnancy records as long as possible ²⁰⁾ . If the information is still missing, imputation by chained equations with the R mice package will be used to address the missing values of aforementioned factors. Datasets will be imputed and results will be summarized over the imputed datasets using Rubin's rules. Number of datasets will be determined on % of pregnancies with missing values.
Subgroup Analyses	Subgroup analysis according to subclass or ingredients of second-generation antipsychotics exposure

Table 11. Sensitivity analyses – rationale, strengths and limitations

	What is being varied? How?	Why? (What do you expect to learn?)	Strengths of the sensitivity analysis compared to the primary	Limitations of the sensitivity analysis compared to the primary
Sensitivity analysis 1	Redefined exposure as having at least one pharmacy dispensing records for antipsychotics after gestational week 4.	To evaluate the effect of exposure misclassification. Women whose estimated antipsychotics exposed window overlapped the pregnancy period but without dispensings after gestational week 4 are potential discontinuers. Non-differential misclassification generally tends to bias results towards the null.	The assumption is that women with at least one dispensing after pregnancy noticeable timing are less likely to be discontinuers.	Reduces sample size of exposed patients
Sensitivity analysis 2	Restricting the maternal psychiatric disorder to schizophrenia only.	To reduce the confounding by indication. First generation antipsychotics are not so often prescribed for mania, bipolar disorder.	Potentially improve exchangeability between the exposure group and comparator.	Reduces sample size of exposed patients, unexposed diseased-comparison group, active comparator group.
Sensitivity analysis 3	Change the lag time between exposure and miscarriage 14days	To evaluate the effect of lag time between exposure and miscarriage. While it is known that there is the lag time between the arrest of development and miscarriage, the lag time between exposure and miscarriage is not completely clear.	Potentially add the knowledge about induction time.	Slightly reduces sample size of unexposed patients.
Sensitivity analysis 4	Exclude initiators during pregnancy in exposed group and active comparator group	To avoid immortal-time bias derived from initiators during pregnancy.	Clarify the exposed mothers as continuers	Reduces sample size of exposed patients

B. Secondary analysis

Hypothesis:	The risk of other selected pregnancy outcomes is elevated with second-generation antipsychotic use during pregnancy.
Exposure contrast:	Second-generation antipsychotics vs Comparator 1: Unexposed, diseased-comparison group Second-generation antipsychotics vs Comparator 2: Active comparator Second-generation antipsychotics vs Comparator 3: Discontinuer
Outcome:	Neonatal outcomes: preterm birth, small for gestational age (SGA), low Apgar score, transfer to NICU, congenital malformations, Maternal outcomes: caesarean section, gestational diabetes, preeclampsia
Analytic software:	R
Model(s):	<p>- <u>Neonatal outcomes : SGA, low Apgar score, transfer to NICU, congenital malformations, preterm birth</u></p> <p>- <u>Maternal outcome : caesarean section, gestational diabetes, preeclampsia</u></p> <p><u>Outcome model:</u> Adjusted RRs will be estimated with modified Poisson regression or binomial regression models in which the second-generation antipsychotics unexposed pregnancies were weighted by the PS distribution of the second-generation antipsychotics exposed pregnancies. We will use robust standard errors to account for correlation within women who participated with >1 pregnancy in this study.</p> <p><u>Propensity score model:</u> logistic regression model, Exposure = covariates</p>
Confounding adjustment method	Propensity scores will be estimated when using logistic regression with exposure as a dependent variable and potential confounders and risk factors for the outcome as independent variables Including all potential confounders and risk factors for the outcome in the propensity score estimation has demonstrated increased precision without increased bias. In the weighted cohort, we will estimate RRs with modified Poisson regression or binomial regression models depending on convergence. Depending on the results of the exploratory data analyses, we may apply other confounding adjustment methods of deemed more appropriate with regard to exposure patterns.
Missing data methods	If information derived from the MBRN is missing, obtain the information from other pregnancy records as long as possible ²⁰ . If the information is still missing, imputation by chained equations with the R mice package will be used to address the missing values of aforementioned factors. Datasets will be imputed and results will be summarized over the imputed datasets using Rubin's rules. Number of datasets will be determined on % of pregnancies with missing values.
Subgroup Analyses	Subgroup analysis according to subclass or ingredients of second-generation antipsychotics exposure

B. Sensitivity analyses in secondary analysis

	What is being varied? How?	Why? (What do you expect to learn?)	Strengths of the sensitivity analysis compared to the primary	Limitations of the sensitivity analysis compared to the primary
Sensitivity analysis 1	Redefined exposure as having at least one pharmacy dispensing records for antipsychotics after gestational week 4.	To evaluate the effect of exposure misclassification. Women whose estimated antipsychotics exposed window overlapped the pregnancy period but without dispensings after gestational week 4 are potential discontinuers. Non-differential misclassification	The assumption is that women with at least one dispensing after pregnancy noticeable timing are less likely to be discontinuers.	Reduces sample size of exposed patients

		generally tends to bias results towards the null.		
Sensitivity analysis 2	Restricting the maternal psychiatric disorder to schizophrenia only.	To reduce the confounding by indication. First generation antipsychotics are not so often prescribed for mania, bipolar disorder.	Potentially improve exchangeability between the exposure group and comparator.	Reduces sample size of exposed patients, unexposed diseased-comparison group, active comparator group.

Other analyses

We will also calculate the estimated excess number of miscarriages by subtracting rates of miscarriage among women with second-generation antipsychotics treated/ first-generation antipsychotics treated /non-treated psychiatric disorders in pregnancy from the rates of miscarriage in the general birth population (per 1000 pregnancies) in Norway. The general pregnancy population will consist of all pregnancies in our linked data file used as the initial population, excluding pregnancies with psychiatric disorders or use of second-generation antipsychotics / first-generation antipsychotics.

7.6. Data sources

7.6.1 Context and rationale for data sources

Reason for selection:

Strengths of data source(s): Our study is based on national health registries with nationwide coverage covering a time period over ten years. The population size allows for analysis on several individual antipsychotics. Important information on maternal characteristics, lifestyle factors and maternal medical complications enables us to control for several important confounding factors. Another important strength with our study is being a prospectively registered nationwide study, which eliminates recall and selection bias. The newly developed pregnancy algorithm to identify early miscarriages and elective terminations (< GW 12) is also one of the strengths⁵⁾. Validity of information on several pregnancy outcomes (preterm birth, birth weight (this relates to SGA), caesarean section, preeclampsia) in MBRN has shown to be very high, which is required to produce robust results^{21–23)}.

Limitations of data source(s):

-Factors leads to less precision: Some potential relevant confounders are unmeasured. These may be substance abuse, poor diet, stress, alcohol use are unmeasured. These may be risk factors for adverse outcomes such as miscarriage²⁴⁾.

Adherence: The exposure measurement is based on dispensed antipsychotic medications in pregnancy. No information about the validity of antipsychotic dispensation is available. However, we expect it to be similar to that of other psychotropics. In particular, antidepressant exposure validity was reported to have a sensitivity of 66.9% and specificity of 99.7%²⁵⁾. Benzodiazepines as anxiolytics and hypnotics exposure validity were reported as a sensitivity of 44.8%, 27.8% and specificity of 99.7%, 100% respectively²⁵⁾.

Data source provenance/curation: The Norwegian data sources are widely used for research and the data holders provide thorough documentation of data contents, assumptions and limitations.

Table 12. Metadata about data sources and software

	Data 1	Data 2	Data 3	Data 4
Data Source(s):	The Medical Birth Registry of Norway (MBRN)	The Norwegian Prescription Registry (NorPD)	The Norwegian Patient Registry (NPR)	Norwegian control and payment of health reimbursements (KUHR)
Study Period:	2004-2018	2004-2019	2008-2019	2006-2019
Eligible Cohort Entry Period:	2008-2018	2008-2018	2008-2018	2008-2018
Data Version (or date of last update):				
Data sampling/extraction criteria:	Personal identification number given to all residents in Norway	Personal identification number given to all residents in Norway	Personal identification number given to all residents in Norway	Personal identification number given to all residents in Norway
Type(s) of data:	Registry	Registry	Registry	Administrative database

Data linkage:	All data			
Conversion to CDM*:	Not reported	Not reported	Not reported	Not reported
Software for data management:				

7.7. Data management

Data files are provided by the Health registries of Norway and labeled according to the project number, version, and date of access. The provided data from these health registries to researchers at UiO are stored in secure servers operated by the University of Oslo called Tjenester for Sensitive Data (TSD). An isolated virtual machine (VM) with a proper operation system and dedicated encrypted storage and CPU resources will be allocated by TSD for each project upon application which stores the received data from registries. The VMs are isolated and only accessible within the UiO network and through secure authentication protocols. Only researchers registered within the specific project number can access TSD and the VM. The project meets the new GDPR requirements as required. Health registries in Norway anonymize data before sending them to TSD by replacing the original IDs with the project-specific generated IDs while preserving the ability to link different data sources. Exporting the results from TSD is strictly limited and needs to be coordinated by the project's admin in a sign-off meeting. Exporting results that can be linked to individual identifications is not permitted. Back-up of the data is automatically and routinely performed multiple times per week within the TSD infrastructure at the University of Oslo.

7.8. Quality control

The data sources have been through extensive quality control procedures by the registry custodian. When new data is received from a registry custodian, the research group has an internal quality check process which includes assessment of reliability and conformance to expected plausible values. Issues are flagged for review by the data quality team and resolved with documentation of decisions made to clean the data before it is released to the research team to conduct studies.

7.9. Study size and feasibility

The source population will include data from the entire Norwegian birthing population between 2008 and 2018, comprising of approximately 860,000 pregnancies, depending on the in- and exclusion criteria that will differ for each objective the study cohorts. Overall antipsychotic exposure prevalence is expected to be 1.16%. Second-generation antipsychotic exposure prevalence is expected to be 0.24%.

Based on an early data extraction (p704), we identified the following approximate numbers of exposed pregnancies:

Analysis	Any antipsychotic	First generation antipsychotic	Second generation antipsychotic
Primary: UiO pregnancy algorithm (2008-2018), exposure in GW0-19	7,000	4,500	2,500
Secondary: MBRN (2008-2018), exposure in pregnancy	6,000	4,500	1,500
Secondary: MBRN (2008-2018), exposure in trimester 1	5,000	3,500	1,500

At a substance level, the following exposures were identified:

ATC code	Antipsychotic	Type	Primary: GW0-19	Secondary: Pregnancy, any time	Secondary: Trimester 1
N05AA01	Chlorpromazine	First generation	600	600	500
N05AA02	Levomepromazine	First generation	500	600	300
N05AB04	Prochlorperazine	First generation	28,00	2,800	2,400
N05AH04	Quetiapine	Second generation	1,700	1,100	900
N05AH03	Olanzapine	Second generation	400	300	300
N05AX12	Aripiprazole	Second generation	300	200	200
N05AX08	Risperidone	Second generation	100	50	50

Table 13. Sample size calculations -cohort study: second-generation antipsychotic exposed vs all unexposed pregnancies (in the whole cohort)

Sample size required to detect associations given 80% study power and type I error rate of 0.05

	RR					RR				
Prevalence of exposure	1.25	1.5	2	3	5	1.25	1.5	2	3	5
	BASELINE RISK of outcome: 10%** MISCARRIAGES					BASELINE RISK of outcome: 8%** PREMATURITY				
0.3%	852000	233333	68000	21333	7000	1092000	300000	88000	28000	9500
1%	255600	70000	20400	6400	2100	327600	90000	26400	8400	2850
2%	127800	35000	10200	3200	1050	163800	45000	13200	4200	1425
5%	51120	14000	4080	1280	420	65520	18000	5280	1680	570
10%	25560	7000	2040	640	210	32760	9000	2640	840	285
	BASELINE RISK of outcome: 5%** LOW BIRTH WEIGHT					BASELINE RISK of outcome: 3%** GESTATIONAL DIABETES/ PRECLAMPSIA				
0.3%	1812000	500000	148000	48000	17000	3092000	855556	254667	83556	30333
1%	543600	150000	44400	14400	5100	927600	256667	76400	25067	9100
2%	271800	75000	22200	7200	2550	463800	128333	38200	12533	4550
5%	108720	30000	8880	2880	1020	185520	51333	15280	5013	1820
10%	54360	15000	4440	1440	510	92760	25667	7640	2507	910
	BASELINE RISK of outcome: 2%** ANY MAJOR CONGENITAL ANOMALY									
0.3%	4692000	1300000	388000	128000	47000					
1%	1407600	390000	116400	38400	14100					
2%	703800	195000	58200	19200	7050					
5%	281520	78000	23280	7680	2820					
10%	140760	39000	11640	3840	1410					

For the primary analysis, 860,000 pregnancies are expected to provide 80% study power to detect a nearly 1.25-fold elevation in miscarriage risk for second-generation antipsychotics compared to other pregnancies in the dataset with 95% confidence intervals excluding the null.

Table 14. Sample size calculations -cohort study: second-generation antipsychotic exposed vs first-generation antipsychotic exposed (in the antipsychotics exposed cohort)

In the feasibility count, there were approximately 7,000 pregnancies with any antipsychotic exposure. The number of second-generation antipsychotic exposure was 2,500 (35%). The number of first-generation antipsychotic exposure was 4,500(65%).

Sample size required to detect associations given 80% study power and type I error rate of 0.05.

	RR					RR				
Prevalence of exposure	1.25	1.5	2	3	5	1.25	1.5	2	3	5
	BASELINE RISK of outcome: 10%** MISCARRIAGES					BASELINE RISK of outcome: 8%** PREMATURITY				
1%	255600	70000	20400	6400	2100	327600	90000	26400	8400	2850
5%	51120	14000	4080	1280	420	65520	18000	5280	1680	570
10%	25560	7000	2040	640	210	32760	9000	2640	840	285
30%	8520	2333	680	213	70	10920	3000	880	280	95
50%	5112	1400	408	128	42	6552	1800	528	168	57
	BASELINE RISK of outcome: 5%** LOW BIRTH WEIGHT					BASELINE RISK of outcome:3%** GESTATIONAL DIABETES/ PRECLAMPSIA				
1%	543600	150000	44400	14400	5100	927600	256667	76400	25067	9100
5%	108720	30000	8880	2880	1020	185520	51333	15280	5013	1820
10%	54360	15000	4440	1440	510	92760	25667	7640	2507	910
30%	18120	5000	1480	480	170	30920	8555	2546	835	303
50%	10872	3000	888	288	102	18552	5133	1528	501	182
	BASELINE RISK of outcome:2%** ANY MAJOR CONGENITAL ANOMALY									
1%	1407600	390000	116400	38400	14100					
5%	281520	78000	23280	7680	2820					
10%	140760	39000	11640	3840	1410					
30%	46920	13000	3880	1280	470					
50%	28152	7800	2328	768	282					

For the primary analysis, 7,000 pregnancies provide 80% study power to detect a nearly 1.5 to 2-fold elevation in miscarriage risk for second-generation antipsychotics compared to first-generation antipsychotics with 95% confidence intervals excluding the null.

8. Limitation of the methods

There are several potential limitations with the methods specified in this protocol.

1. The data were not collected for research and some important variables may not be collected or will be measured imperfectly
 - a. We have selected validated algorithms when possible
 - b. We have created proxies for important variables that are not directly captured in the data to reduce confounding by unmeasured factors
2. There will not be randomization
 - a. We will balance exposure groups on important risk factors for the exposure and outcome(s)

9. Protection of human subjects

The proposed study is observational research that makes secondary use of data collected as part of routine care and does not involve any intervention, alteration in standard clinical care or use of any procedure in patients. Therefore, there will be no adverse events related to the study itself. No patients will be contacted for any of the proposed studies. Prior to our acquisition of the data, all personal identifiers will be encrypted. This encryption minimizes the risk of patient reidentification in the unlikely event of a breach in data security.

10. Reporting of adverse events

The proposed study is observational research that makes secondary use of data collected as part of routine care and does not involve any intervention or alteration in clinical care. Therefore, reporting of adverse events related to this study is not applicable. Safety evaluations for this study are limited to the specified safety outcomes stated in section 4.4.2.

11. References

- 1) Reutfors J, Cesta CE, Cohen JM, et al. Antipsychotic drug use in pregnancy: A multinational study from ten countries. *Schizophr Res.* 2020 Jun;220:106-115. PMID: 32295750.
- 2) Huybrechts KF, Straub L, Karlsson P, et al. Association of In Utero Antipsychotic Medication Exposure With Risk of Congenital Malformations in Nordic Countries and the US. *JAMA Psychiatry.* 2023 Feb 1;80(2):156-166. PMID: 36477338.
- 3) Liu X, Kolding L, Momen N, et al. Maternal antipsychotic use during pregnancy and congenital malformations. *Am J Obstet Gynecol MFM.* 2023 Jun;5(6):100950. PMID: 37015311.

- 4) Sørensen MJ, Kjaersgaard MI, Pedersen HS, et al. Risk of Fetal Death after Treatment with Antipsychotic Medications during Pregnancy. *PLoS One*. 2015 Jul 10;10(7):e0132280. PMID: 26162087.
- 5) Nordeng H, Lupattelli A, Engjom HM, van Gelder MMHJ. Detecting and Dating Early Non-live Pregnancy Outcomes: Generation of a Novel Pregnancy Algorithm From Norwegian Linked Health Registries. *Pharmacoepidemiol Drug Saf*. 2024 Sep;33(9):e70002. PMID: 39238438.
- 6) Tang W, Zhou LJ, Zhang WQ, et al. Adverse perinatal pregnancy outcomes in women with schizophrenia: A systematic review and meta-analysis. *Schizophr Res*. 2023 Dec;262:156-167. PMID: 37979419.
- 7) Jarde A, Morais M, Kingston D, et al. Neonatal Outcomes in Women With Untreated Antenatal Depression Compared With Women Without Depression: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. 2016 Aug 1;73(8):826-37. PMID: 27276520.
- 8) Grigoriadis S, VonderPorten EH, Mamisashvili L, et al. The impact of maternal depression during pregnancy on perinatal outcomes: a systematic review and meta-analysis. *J Clin Psychiatry*. 2013 Apr;74(4):e321-41. PMID: 23656857.
- 9) Rusner M, Berg M, Begley C. Bipolar disorder in pregnancy and childbirth: a systematic review of outcomes. *BMC Pregnancy Childbirth*. 2016 Oct 28;16(1):331. PMID: 27793111.
- 10) Fabre C, Pauly V, Baumstarck K, et al. Pregnancy, delivery and neonatal complications in women with schizophrenia: a national population-based cohort study. *Lancet Reg Health Eur*. 2021 Sep 7;10:100209. PMID: 34806069.
- 11) Edvardsson K, Hughes E, Copnell B, et al. Severe mental illness and pregnancy outcomes in Australia. A population-based study of 595 792 singleton births 2009-2016. *PLoS One*. 2022 Feb 28;17(2):e0264512. PMID: 35226688.
- 12) Magnus MC, Havdahl A, Morken NH, et al. Risk of miscarriage in women with psychiatric disorders. *Br J Psychiatry*. 2021 Sep;219(3):501-506. PMID: 33448259
- 13) Sakai T, Ohtsu F, Mori C, et al. Signal of Miscarriage with Aripiprazole: A Disproportionality Analysis of the Japanese Adverse Drug Event Report Database. *Drug Saf*. 2017 Nov;40(11):1141-1146. PMID: 28664356.
- 14) Twaites BR, Wilton LV, Shakir SA. The safety of quetiapine: results of a post-marketing surveillance study on 1728 patients in England. *J Psychopharmacol*. 2007 Jun;21(4):392-9. PMID: 17656426.
- 15) Goldstein DJ, Corbin LA, Fung MC. Olanzapine-exposed pregnancies and lactation: early experience. *J Clin Psychopharmacol*. 2000 Aug;20(4):399-403. PMID: 10917399.
- 16) Coppola D, Russo LJ, Kwart RF Jr, et al. Evaluating the postmarketing experience of risperidone use during pregnancy: pregnancy and neonatal outcomes. *Drug Saf*. 2007;30(3):247-64. PMID: 17343431.
- 17) Irgens LM. The Medical Birth Registry of Norway. Epidemiological research and surveillance throughout 30 years. *Acta Obstet Gynecol Scand*. 2000 Jun;79(6):435-9. PMID: 10857866.

- 18) Sundermann AC, Mukherjee S, Wu P, et al. Gestational Age at Arrest of Development: An Alternative Approach for Assigning Time at Risk in Studies of Time-Varying Exposures and Miscarriage. *Am J Epidemiol*. 2019 Mar 1;188(3):570-578. PMID: 30521025.
- 19) Lau B, Cole SR, Gange SJ. Competing risk regression models for epidemiologic data. *Am J Epidemiol*. 2009 Jul 15;170(2):244-56. PMID: 1949424
- 20) Sakai T, Nordeng H, van Gelder MMHJ. Validity of methods to infer missing data on maternal characteristics using a medical birth registry: longitudinal methods versus multiple imputation. [Under review]
- 21) Lehmann S, Baghestan E, Bør Dahl P, et al. Validation of data in the Medical Birth Registry of Norway on delivery after a previous cesarean section. *Acta Obstet Gynecol Scand*. 2017;96(7):892-897. PMID: 28196281.
- 22) Moth FN, Sebastian TR, Horn J, et al. Validity of a selection of pregnancy complications in the Medical Birth Registry of Norway. *Acta Obstet Gynecol Scand*. 2016;95(5):519-527. PMID: 26867143.
- 23) Thomsen LC, Klungsøyr K, Roten LT, et al. Validity of the diagnosis of preeclampsia in the Medical Birth Registry of Norway. *Acta Obstet Gynecol Scand* 2013;92(8):943-950. PMID: 23621424.
- 24) Quenby S, Gallos ID, Dhillon-Smith RK, et al. Coomarasamy A. Miscarriage matters: the epidemiological, physical, psychological, and economic costs of early pregnancy loss. *Lancet*. 2021 May 1;397(10285):1658-1667. PMID: 33915094.
- 25) Skurtveit S, Selmer, R, Odsbu I, et al. Self-reported data on medicine use in the Norwegian Mother and Child cohort study compared to data from the Norwegian Prescription Database. *Norsk Epidemiologi*, 2014, 24(1-2).

12. Appendices

12.1.1. Appendix E. Tentative flow chart

